Peak and Range of Blood Glucose are also associated with ICU Mortality

Geoffrey M. Shaw¹, J. Geoffrey Chase², Dominic Lee³, Maxim Bloomfield⁴, Carmen V. Doran², Jessica Lin² and Thomas Lotz²

¹ Canterbury District Health Board, Dept of Intensive Care and University of Otago, Christchurch School of Medicine and Health Sciences, Christchurch, NZ

² Centre for Bioengineering, Dept of Mechanical Engineering, University of Canterbury, Christchurch, NZ

³ Dept of Mathematics and Statistics, University of Canterbury, Christchurch, NZ

⁴ University of Otago School of Medicine and Health Sciences, Christchurch, NZ

The main contributing factor for the reduction in mortality of critically ill patients under tight glycemic control is debated between tight control of glucose levels [1, 2] and the administration of insulin [3]. This study determines the relationships between several blood glucose markers and mortality in the critically ill.

Hypothesis:

This study has two main hypotheses:

- Maximum blood glucose level is a marker for ICU mortality.
- Maximum range of blood glucose values during stay is a marker for ICU mortality.

Outline of Study

Plasma glucose values, insulin infusions, age, sex, APACHE II score, length of stay, ICU mortality, mechanical ventilation status and primary diagnosis were recorded for patients with a stay >72 hours in the Christchurch ICU.

Patient Cohort

A total of 1164 patients were admitted to the Christchurch ICU during the 2003. Of these patients, 241 had a stay greater than 3.0 days. Only 201 who had complete plasma glucose data were included in the retrospective data audit.

Christchurch Hospital Intensive Care Unit Patient Subgroups

Subgroup	Number of Patients	% of total
Cardiac	32	16%
Pulmonary	34	17%
Sepsis	15	7%
Other Medical	26	13%
Neurologic	19	9%
General Surgical	41	20%
Trauma	34	17%
TOTAL	201	100%

The patients were divided into the categories used by Krinsley [5] depending on their primary diagnosis. This subset is similar to that in Krinsley, with the major difference being the size of the populations examined.

General Description of Patients in Christchurch Hospital Intensive Care Unit

Subgroup	Number of Patients	Age (years) ^a	Female	APACHE II ^b	Mech. Vent. ^c	LOS (days) ^d	ICU Survival ^e
Cardiac	32	65.3 (30-85)	31%	22.1 (8-39)	78%	6.4 (3.0-21.8)	94%
Pulmonary	34	59.0 (18-82)	56%	20.5 (15-36)	38%	11.9 (3.1-36.0)	71%
Sepsis	15	54.4 (1-82)	47%	20.3 (15-36)	60%	12.0 (3.1-36.0)	93%
Other Medical	26	54.2 (20-84)	58%	20.4 (17-36)	54%	10.4 (3.0-39.9)	88%
Neurologic	19	54.1 (1-81)	42%	18.1 (7-24)	74%	8.5 (3.0-35.0)	84%
General Surgical	41	64.6 (17-85)	32%	21.5 (10-41)	68%	10.6 (3.0-42.5)	88%
Trauma	34	34.9 (16-83)	16%	17.2 (8-30)	55%	10.7 (3.0-60.0)	94%
TOTAL	201	55.6 (1-85)	39%	21.8 (7-45)	66%	9.9 (3.0-60.0)	87%

^a Age, APACHE II score and LOS are expressed as mean (range)
^b Acute Physiology and Chronic Health Evaluation score II
^c Mechanical Ventilation
^d Length of Stay (in increments of 0.1)
^e Intensive Care Unit Survival (obtained from ICU discharge records)

Patient Parameters in Survivors and Non-survivors in Christchurch Hospital Intensive Care Unit

Subgroup	Number of Patients	Age (years)	APACHE II ^b	Mech. Vent. ^c	LOS (days) ^d
All	27	56.3 (10-82)	26.4 (7-41)	74%	11.3 (3.1-36.0)
Survivors	174	54.3 (1-85)	21.4 (8-36)	66%	9.8 (3.0-60.0)
Cardiac	2	57.5 (56-60)	31.0 (23-39)	100%	3.5 (3.1-3.8)
Survivors	30	55.8 (30-85)	21.4 (8-36)	66%	6.6 (3.0-21.8)
Pulmonary	10	63.7 (24-82)	22.0 (20-26)	50%	13.0 (4.7-36)
Survivors	24	57.0 (18-79)	26.5 (15-36)	33%	10.8 (3.1-24.8)
Sepsis	1	48.0	27.0	100%	12.8
Survivors	14	55.9 (1-82)	26.2 (21-36)	57%	11.9 (3.0-39.9)
Other Medical	4	66.0 (47-84)	27.7 (25-33)	75%	9.0 (3.4-22.3)
Survivors	22	52.0 (20-84)	25.2 (17-36)	50%	10.8 (3.0-35.0)
Neurologic	3	64.3 (53-78)	15.7 (7-20)	67%	5.6 (3.5-7.1)
Survivors	16	52.1 (1-81)	18.7 (11-24)	75%	9.1 (3.2-43.8)
General Surgical	5	69.2 (59-77)	27.0 (18-41)	80%	16.1 (3.6-32.3)
Survivors	36	63.9 (17-85)	20.4 (10-35)	67%	9.8 (3.0-42.5)
Trauma	2	23.0 (20-26)	25.0	100%	6.5 (4.5-8.4)
Survivors	32	32.2 (15-83)	16.8 (8-30)	84%	11.0 (3.0-60.0)

^a Age, APACHE II score and LOS are expressed as mean (range)
^b Acute Physiology and Chronic Health Evaluation score II
^c Mechanical Ventilation
^d Length of Stay

There was no significant difference in age, APACHE II score, length of stay or mechanical ventilation rate between survivors and non-survivors using a P < 0.05 statistical significance value. Generally, there was a trend towards younger survivors in the cardiac (P = 0.15) and pulmonary (P = 0.13) groups, as well as across all categories (P = 0.13). Survivors also tended to have a lower APACHE II score across all categories (P = 0.09). To better confirm significance in these subgroups, a larger cohort would be required.

Associated risk of death (ROD) is shown in each category. The distribution is relatively normal, with 26% of patients studied having an APACHE II score in the range of 20 - 24.

Patients with a stay of 3 - 4 days made up 19.1% of the patient population studied, while patients with 3 - 9 days stay made up 62.4% of the population. The longest length of stay in the ICU during 2003 was 60.0 days.

Of 3652 blood glucose measurements over all patients, the majority (34%) were 6 - 8 mmol/L. There were 116 (3%) hypoglycemic values recorded by 81 patients (hypoglycemic threshold is set at 4 mmol/L).

Conclusions

Mortality increased with increasing maximum blood glucose, range of blood glucose and trapezoidal mean of blood glucose, suggesting that all three measures have a negative effect on the survival of an ICU patient. Equally importantly, these three measures were seen to be markers of increased mortality independent of APACHE II scores. From the results, it is proposed:

- Maximum blood glucose level, and hence the range of blood glucose levels, are also significant factors associated with mortality, in addition to the mean blood glucose value.
- Frequent measurement and tight control need to be applied immediately following ICU admission for effective blood glucose management.
- Both absolute blood glucose levels and control of blood glucose variation are important factors when considering new approaches to tight glycemic control in critical illness.

Hence, this retrospective study clearly provides additional metrics for considering blood glucose control and illustrates the difficulty in controlling blood glucose levels in intensive care. **Significant improvements in the measurement and practice of treating hyperglycaemia will need to be developed to fully realize the benefits of tight control.**

- Van den Berghe, G. Wouters, P. Weekers, F. Verwaest, C. Bruyninckx, F. Schetz, M. Vlasselaers, D. Ferdinande, P. Lauwers, P. Bouillon, R. Intensive insulin therapy in the critically ill patients. *N Engl J Med* 2001; 345(19):1359-1367
- Van den Berghe, G. Wouters, P.J. Bouillon, R. Weekers, F. Verwaest, C. Schetz, M. Vlasselaers, D. Ferdinande, P. Lauwers, P. Outcome benefit of intensive insulin therapy in the critically ill: Insulin dose versus glycemic control. *Crit Care Med* 2003; 31(2):359-366
- Anname, D. Melchior, J.C. Hormone replacement therapy for the critically ill. *Crit Care Med* 2003; 31(2):634-635.
- Burden, R. and J. Faires, Numerical Analysis. Fourth Edition. PWS-KENT Publishing Company, 1989
- Krinsley, J. S. 2003. Association between Hyperglycemia and Increased Hospital Mortality in a Heterogeneous Population of Critically Ill Patients. *Mayo Clin Proc.* 78(12): 1471-1478

Calculating the Blood Glucose Mean

Calculation of mean blood glucose values for irregularly sampled data required the use of a random walk to bridge the gaps between data and trapezoidal assumptions to determine confidence intervals for the mean.

Assuming that the blood glucose readings are ergodic, and using the trapezoidal rule outlined in Burden and Faires [4], an estimate of the patient's mean blood glucose for the duration [0, T] is defined and illustrated.

Using the trapezoidal mean, it is assumed that if blood glucose had been measured between two measured points, then it would lie on the linear line between those two points. However, this assumption may not hold. Therefore, a method for estimating the range within which this intermediate blood glucose measurement lies is required.

A 90% confidence interval can be determined:

To put the mean glucose values into context, confidence intervals were calculated. However, it was found that the length of the intervals was far too long to be helpful, due to infrequent sampling. The linear least squares fit shown in Figure 2.10 considers only patients with a half-bandwidth less than 6 mmol/L, and consequently a confidence interval length of less than 12 mmol/L. This 12 mmol/L value represents an extreme size for a confidence interval and would render the associated mean value useless.

Therefore blood glucose must be measured every hour or more frequently to ensure a confidence interval of ±1mmol/L or less on the mean value.

Results

Effect of Frequency, Maximum and Range of Blood glucose Measurements on Survival in Christchurch Hospital Intensive Care Unit

Subgroup	Number of Patients	Average Measurements per day ^a	Maximum Blood Glucose (mmol/L) ^b	Blood Glucose Range (mmol/L) ^{c,d}
All	27	4.7 (0.2-13.0)	14.1 (7.1-22.2)	8.4 (1.5-15.7)
Survivors	174	3.1 (0.1-16.2)	10.8 (3.9-26.9)	6.1 (0-26.7)
Cardiac	2	10.3 (8.8-11.9)	15.9 (14.7-17.1)	12.6 (9.7-15.4)
Survivors	30	4.4 (0.2-9.8)	12.1 (8.2-19.8)	7.6 (2.2-18.0)
Pulmonary	10	4.3 (0.4-8.0)	12.0 (6.9-18.3)	7.8 (1.5-13.3)
Survivors	24	3.2 (0.1-7.9)	11.1 (6.0-17.6)	6.5 (0.0-14.8)
Sepsis	1	7.2	11.5	9.7
Survivors	14	3.5 (0.1-8.0)	11.5 (4.8-17.6)	7.4 (0.0-14.3)
Other Medical	4	7.9 (1.4-13.0)	16.6 (12.9-22.2)	12.9 (8.1-16.7)
Survivors	22	4.7 (0.1-11.2)	14.3 (6.0-28.9)	9.5 (0.0-26.7)
Neurologic	3	4.4 (0.5-7.2)	13.7 (7.2-18.4)	8.5 (1.7-11.7)
Survivors	16	2.4 (0.1-10.0)	9.2 (3.5-18.4)	3.9 (0.0-14.3)
General Surgical	5	2.0 (0.2-5.0)	11.1 (7.8-13.8)	6.4 (3.0-10.1)
Survivors	36	2.7 (0.1-10.3)	10.0 (5.4-19.9)	5.4 (0.0-14.7)
Trauma	2	0.8 (0.6-0.7)	13.8 (12.4-15.4)	7.4 (5.1-9.7)
Survivors	32	1.2 (0.1-16.2)	8.3 (4.5-16.6)	3.1 (0.0-13.9)

^a Average Measurements per day. Maximum Blood Glucose and Blood Glucose Range are expressed as mean (range)
^b Calculated by dividing total number of measurements by total length of stay
^c Maximum recorded blood glucose measurement during stay
^d Maximum recorded minus minimum recorded blood glucose measurements during stay

The maximum value of blood glucose along with the range of blood glucose values was higher in non-survivors (P = 0.001 and P = 0.003, respectively), suggesting that the level of deviation from basal blood glucose values has a negative impact on survival in critically ill patients, especially in trauma patients (P = 0.04 and P = 0.05, respectively). The maximum values of blood glucose correlate well with those found in Krinsley [5] for both survivors (10.8 mmol/L compared to 9.8 mmol/L) and non-survivors (14.1 mmol/L compared to 14.3 mmol/L).

Effect of Blood Glucose Trapezoidal Means of the 86 Patients on Insulin on Survival in Christchurch Hospital Intensive Care Unit

Subgroup	Number of Patients	Trapezoidal Mean Blood Glucose (mmol/L) ^a
All	16	9.3 (8.5-14.1)
Survivors	70	7.9 (4.5-12.2)
Cardiac	2	9.8 (8.5-10.1)
Survivors	12	7.5 (6.9-9.8)
Pulmonary	6	8.0 (7.0-9.8)
Survivors	12	7.6 (6.9-9.8)
Sepsis	1	7.4
Survivors	8	8.2 (6.1-10.0)
Other Medical	3	10.7 (8.5-14.1)
Survivors	13	8.4 (6.9-10.0)
Neurologic	2	11.6 (10.2-13.0)
Survivors	4	9.1 (7.2-12.2)
General Surgical	2	9.3 (8.1-9.9)
Survivors	11	7.5 (6.5-9.8)
Trauma	-	-
Survivors	1	9.7

^a Trapezoidal mean calculated using Equation 2, expressed as mean (range)

Maximum Blood Glucose Value and Intensive Care Unit Survival in Christchurch Hospital Intensive Care Unit

Maximum Blood Glucose (mmol/L)	Number of Patients	ICU Survival ^a
<6	14	100%
6-8	27	92%
8-10	48	96%
10-12	34	85%
12-14	30	70%
14-16	27	79%
16-18	9	78%
18-20	6	87%
>20	6	83%

^a Maximum recorded blood glucose measurement during stay
^b Intensive Care Unit Survival

Blood Glucose Trapezoidal Mean and Intensive Care Unit Survival in Christchurch Hospital Intensive Care Unit

Blood Glucose Trapezoidal Mean (mmol/L)	Number of Patients	ICU Survival ^a
<6	4	100%
6-8	16	88%
8-10	32	88%
10-12	19	80%
12-14	19	84%
14-16	7	29%

^a Trapezoidal mean calculated using Equation 2
^b Intensive Care Unit Survival

Effect of Insulin Infusion on Survival in Christchurch Hospital Intensive Care Unit

Subgroup	Number of Patients	Average Insulin Infused (U/hr) ^a	Proportion of Stay on Insulin ^b
All	27	1.1 (0-5.7)	41% (0-97)
Survivors	174	0.5 (0-5.7)	24% (0-88)
Cardiac	2	3.8 (3.4-3.8)	90% (84-97)
Survivors	30	0.6 (0-3.7)	41% (0-88)
Pulmonary	10	0.7 (0-2.1)	41% (0-79)
Survivors	24	0.4 (0-2.4)	27% (0-84)
Sepsis	1	1.2	89%
Survivors	14	0.5 (0-1.9)	27% (0-83)
Other Medical	4	2.6 (0-5.7)	67% (0-83)
Survivors	22	1.0 (0-5.7)	43% (0-97)
Neurologic	3	1.0 (0-1.8)	45% (0-74)
Survivors	16	0.4 (0-3.2)	16% (0-80)
General Surgical	5	0.2 (0-1.0)	13% (0-81)
Survivors	36	0.3 (0-1.9)	17% (0-80)
Trauma	2	0.0	0%
Survivors	32	0.1 (0-1.6)	2% (0-47)

^a Average insulin infused. Proportion of stay on insulin are expressed as mean (range)
^b Calculated by dividing total insulin infused by length of time between first and last infusions
^c Calculated by dividing number of hours on insulin divided total length of stay in hours

Patients with a maximum recorded blood glucose value less than 6 mmol/L had the highest survival rate. The trend shows that as the maximum blood glucose increases, the chance of ICU survival decreases. Similar trends were seen for the range and trapezoidal mean of blood glucose