THESIS

FOR

MASTER OF SCIENCE AND HONOURS

University of New Zealand

1938

-----oo000-----

A.J.C. Nicholson

Codeword:

PHYSICAL
SCIENCES
LIBRARY
THESIS

copy 2

THE THERMODYNAMIC DISSOCIATION CONSTANTS
OF OXALIC ACID IN WATER AND
METHANOL-WATER MIXTURES

INDEX

	Page
INTRODUCTION,	1.
EXPERIMENTAL METHOD.	
Materials,	20.
Electrodes,	24.
Cells,	25.
CALCULATION OF RESULTS.	
First Dissociation Constant,	29.
Second Dissociation Constant,	36.
Tables,	37.
Graphs,	38.
DISCUSSION,	44 .
SUMMARY,	53.
REFERENCES.	54.

INTRODUCTION

In the classical theory of acid-base dissociation acids and bases were characterised by their ability to split off hydrogen ions and hydroxyl ions respectively. This definition does not explain the complementary but opposite properties of acids and bases, nor account for their behaviour in non-aqueous media. A more satisfactory definition is that suggested in 1923 by Bronsted¹ and Lowry² independently, in the scheme : -

according to which an acid is a substance which can donate a proton, thus forming a base, while conversely, a base can accept a proton to form an acid.

An acid, therefore, always has one more positive charge than a base, and the following are typical examples of various types of acids.

This theoretical scheme must be modified in practice by the fact that the proton cannot exist in a

free state, but can only be transferred from one molecule to another. The solvent molecules must be regarded under this definition as acids and bases, giving : -

where A_1 and B_1 represent a corresponding acid-base pair. This 'protolysis' is seen to be a general reaction including the 'electrolytic dissociation' of acids and bases, 'hydrolysis' of salts, 'neutralisation' as treated in the classical theory, as special cases. In aqueous solutions water can play the part of either an acid or a base, i.e.,

The strength of an acid can most readily be measured by the size of the dissociation constant

$$K = \frac{[\text{H}^+][\text{B}]}{[\text{A}]}$$

where the square brackets denote various types of concentration expressions as discussed later.

For a dibasic acid, H_2A , two such constants are

obtained:

For a dibasic organic acid the comparison of the first and second dissociation constants gives some information about the molecular structure. For a symmetrical acid the two carboxyl groups will show equal tendencies to dissociate. In this case K_1 should be four times as great as K_2 on statistical grounds alone; for the acid $\text{R}(\text{COOH})_2$ has twice as many ionisable hydrogen atoms as $\text{R}(\text{COOH})\text{COO}^-$, while in addition the anion $\text{R}(\text{COO})_2^-$ will have twice as great a tendency to unite with hydrogen ion as the anion $\text{R}(\text{COOH})\text{COO}^-$. A comparison of K_2 with $\frac{K_1}{4}$ therefore measures the effect of the dissociation of one carboxyl group on the other. The effect of the acquired charge on one carboxyl may be transmitted through the molecule by displacement of electrons (inductive effect), or directly as a pure electrostatic effect through the medium. Considering this second effect only, for a symmetrical acid, Bjerrum⁴⁴ has calculated from Boltzmann's potential energy concentration relationship that

$$\log \frac{K_1}{K_2} - \log 4 = \frac{3.1 \times 10^{-8}}{\gamma}$$

where γ is the distance between the ionizing groups. This

gives values of $\bar{v}$ of the right order for intramolecular distances, but for a relatively strong acid, such as oxalic acid in its first dissociation, the values are low, and the inductive effect, and also the effect of the field of the pole on hydrogen ion concentration, must be allowed for. Gane and Ingold¹⁴ deduce the following theoretical formula allowing for these factors: -

$$\log \frac{K_1}{K_2} - \log 4 = \frac{q\phi}{2.3026kT}$$

where q = the charge on the ion, k = the Boltzmann constant, and T = the absolute temperature. The value of $\bar{v}$ is then calculated by the further expression for 25°C: -

$$\phi = \frac{6.112 \times 10^{-12}}{\bar{v}} + \frac{1.368 \times 10^{-41}}{\bar{v}^5}$$

These values for $\bar{v}$ are higher for oxalic acid and nearer the values for intramolecular distances obtained by other methods.

The general theoretical effect of solvent on acid-base dissociation has been discussed by Bronsted³, Richardson⁴, and others. The strength of a weak acid depends largely on the extent of its dissociation, which depends to a large extent on the electrostatic force between the ions, and is hence affected by the dielectric constant.

For acids of 4 general types : -

the products of dissociation attract one another in the first two cases, have no action in the third, and repel one another in the fourth. It follows that lowering of the dielectric constant (as by transfer from aqueous to alcoholic solution) will diminish the strength in the first two cases, have no effect in the third, and increase it in the fourth.

A more quantitative treatment after Bronsted can be derived as follows: - ⁴⁵

The classical electrostatical formula for the change in energy of an ion on transfer from one solvent to another of different dielectric constant is : -⁴⁶

$$\frac{q^2}{2a} \left(\frac{1}{D_{H_2O}} - \frac{1}{D_s} \right)$$

where q = the charge on the ion, a its radius, D_{H_2O} the dielectric constant in one solvent and D_s in the other, these latter in the present case being water and alcohol water mixtures.

Consider an electrolyte with the ions present in equilibrium in each of the two solvents.

Starting with a gram-mol of undissociated acid in water at unit concentration, produce a gram-mol of each ion at unit concentration by a reversible process (i.e. Van't Hoff equilibrium box).

$$\text{Then work done} = RT \ln K_{H_2O}$$

Transfer these separate ions to the other solvent at unit concentration and the energy change is

$$2 \times \frac{Nq^2}{2a} \left(\frac{1}{D_{H_2O}} - \frac{1}{D_s} \right)$$

where N = Avagadro number.

The ions are now transferred to the undissociated state at unit concentration in this solvent, and the work term is : -

$$RT \ln K_s$$

If we make the reasonable assumption that undissociated molecules can be transferred from one solvent to the other without work, the cycle is complete

$$\text{and} \quad \sum \text{work} = 0$$

$$\text{i.e.} \quad RT \ln K_{H_2O} - RT \ln K_s = \frac{2Nq^2}{2a} \left(\frac{1}{D_{H_2O}} - \frac{1}{D_s} \right)$$

$$\text{or} \quad pK_s - pK_{H_2O} = \frac{Nq^2}{2 \cdot 3026 a RT} \left(\frac{1}{D_{H_2O}} - \frac{1}{D_s} \right)$$

For a dibasic acid in its second stage of dissociation : -

The dielectric work consists of three terms: -

$$\text{for } \text{H}^+, \quad \frac{Nq^2}{2a} \left(\frac{1}{D_{\text{H}_2\text{O}}} - \frac{1}{D_s} \right)$$

$$\text{for } \text{X}^{=} \quad \frac{N4q^2}{2a} \left(\frac{1}{D_{\text{H}_2\text{O}}} - \frac{1}{D_s} \right)$$

$$\text{and for } \text{HX}^- \quad \frac{Nq^2}{2a} \left(\frac{1}{D_{\text{H}_2\text{O}}} - \frac{1}{D_s} \right)$$

$$\text{giving} \quad \frac{2 \times N \times 2q^2}{2a} \left(\frac{1}{D_{\text{H}_2\text{O}}} - \frac{1}{D_s} \right)$$

$$\text{and } \text{pK}_s - \text{pK}_w = \frac{2 Nq^2}{2 \cdot 3026 a RT} \left(\frac{1}{D_{\text{H}_2\text{O}}} - \frac{1}{D_s} \right)$$

From these results a value of 'a' can be calculated or possibly, if 'a' does not remain constant, a mean of the values of 'a' in the two solvents.

In the present work, the dissociation constants of oxalic acid are studied: -

In both these cases, lowering of the dielectric

constant should diminish the strength of the acid. Since there is a greater force of attraction between the doubly charged ion and the proton in the second dissociation, the effect of change in solvent should be more marked here than in the first case. Richardson⁴ considers from his studies of zwitterions that the charges of divalent ions tend to function almost as discrete charges, which may make the change in dissociation of the divalent acid less than anticipated.

An increase in strength is also caused by an actual chemical affinity between solvent and the dissociated ions. In the same way that protons combine with water to give oxonium ions, they will combine with other hydroxylic solvents such as alcohol, and the dissociation of an acid may be regarded as a substitution in which solvent molecules compete with one ion for association with the other. Bronsted³ considers that the dielectric effect is the predominating one, and while this may not be the case with acetone and other solvents, in pure methanol the solvation will be high and of a similar type to that of water. Hartley, Butler and others,⁵ consider that in mixed aqueous solvents ions tend to associate with water rather than the other solvent present, and, so long as there is sufficient water in the mixture, they behave almost as though in aqueous solution. For instance, these workers showed that Lithium ions behave in certain respects almost identically in water, and a 30%

alcohol water mixture, and that the addition of as little as 1% water into a non-hydroxylic solvent causes a marked change in the dissociation. Harned and Thomas⁶ have shown that the mean distance of approach of the ions for acetic acid is the same in water, 10% methanol-water, and 20% methanol-water mixtures. These solvents were used in the present work, hence the effect of change of solvent can be regarded very largely as due to the change in dielectric constant.

The methods used up to the present for determining the dissociation constants of acids can be grouped under the following four headings.

The conductivity method as originally developed by Ostwald, Walden, Voermann, and many others. Using the notation of Brønsted, this gives a 'constant' K_c expressed in terms of the concentrations of the reactants, which varies considerably with dilution. It is assumed that the 'Arrhenius coefficient' $\left(\frac{\Lambda_0}{\Lambda_\infty} \right)$ gives the degree of ionic dissociation, which is certainly not strictly accurate. Using this method, Chandler⁹ obtained the values .038, and 4.9×10^{-5} , for the first and second dissociation constants, respectively, of oxalic acid, and Brucker¹¹ .057, and 6.9×10^{-5} .

A second method consists in measuring the number of

hydrogen ions present by their catalytic effect on the velocity of any suitable reaction. There is still considerable doubt as to what constant this method gives, but it would presumably be K_c as in the conductivity method. Glasstone⁴³ considers the evidence is inconclusive as to whether catalytic power is proportional to concentrations or to activities. Using these methods and molar concentrations, the following values are found : - first dissociation constant $\cdot 040$, Hornel and Butler⁴⁹ $\cdot 057$; second dissociation constant $7\cdot 2 \times 10^{-5}$, Dawson and Smith,¹⁰.

A third method used by many workers is that of electrometric titration. The calomel electrode is used, and the liquid junction is 'annulled' by means of various types of salt-bridge. This gives a constant which, while better than K_c , also varies considerably with dilution. Very large errors arise in this method from uncertainty with regard to the 'annulled' liquid junction potentials. In 1936, Hamer and Acree¹² made determinations of the errors in pH, dissociation constants, ionic strength, and hydrogen ion activity coefficients, arising from neglect of corrections, or from applications of partial corrections, for the potential of the liquid junction between saturated KCl and sodium malonate buffer solutions. They showed that partial corrections produce a larger error in these quantities than no correction at all, and that an error as small as $\cdot 3\%$ in pH may produce a change as large as 23% in dissociation constants.

The wide range of values by this method is shown in the following results: - first dissociation constant: $\cdot 035$ Enklaar¹³; $\cdot 059$ Gane and Ingold¹⁴; $\cdot 17$ Britton¹⁵; second dissociation constant: $4\cdot 7 \times 10^{-5}$ Enklaar¹³; $6\cdot 4 \times 10^{-5}$ Gane and Ingold¹⁴; $13\cdot 8 \times 10^{-5}$ Britton¹⁵.

Other values cited for oxalic acid without any indication of which constant is meant are $\cdot 063$, and $7\cdot 94 \times 10^{-5}$, Richardson⁴; $\cdot 02$, and 3×10^{-5} , Glasstone¹⁶; 5×10^{-5} , Philbrick¹⁷.

A fourth method has been recently evolved by Harned and Owen¹⁸, and modified by Harned and Ehlers¹⁹, which uses cells without liquid junctions. This gives a thermodynamic dissociation constant

$$K_a = \frac{a_+ a_-}{a}$$

which is independent of the dilution.

A considerable number of acids have been studied by this method and results obtained in excellent agreement with independent methods. Harned and Ehlers¹⁹ obtained $K = 1\cdot 754 \times 10^{-5}$ for acetic acid by this method, and MacInnes and Shedlovsky²⁰ $K = 1\cdot 753 \times 10^{-5}$ by conductivity measurements, corrected for ionic attraction by means of the Debye-Huckel-Onsager equations.

Using the same method as modified for the second

dissociation constant of an acid, (as also used in the present work), Nims²¹ determined the second dissociation constant of phosphoric acid. His values are in good agreement with values determined by Bjerrum and Unmack²² using an electrometric titration method with a calomel electrode.

Other acids measured by this method are formic,²³ 1.77×10^{-4} ; propionic²⁴, 1.34×10^{-5} ; butyric²⁵, 1.51×10^{-5} ; chloracetic²⁶, 1.38×10^{-3} ; glycollic²⁷, 1.48×10^{-4} ; 1st phosphoric²⁸, 7.52×10^{-3} ; 2nd sulphuric²⁹, 1.2×10^{-2} .

All these acids are much weaker in their first dissociation constant than oxalic acid, but their values are about the same order as that of the second dissociation constant for oxalic.

Very little accurate work has been done on dissociation constants in solvents other than water. Harned and Ernbee³⁰ have measured the dissociation constant of acetic acid in 10% and 20% methanol-water mixtures.

A previous investigator in this laboratory has determined the dissociation constants by this method, and obtained the values $\cdot 01718$ and 5.012×10^{-5} at 25°C . The first result will be shown to be erroneous, but the errors made do not affect the second dissociation constant, and this value is used in the present work for comparison purposes.

In the present work, the method used by Harned and

his co-workers is employed to measure the thermodynamic first dissociation constant of oxalic acid at 25°C, 30°C, and 35°C, and the first and second dissociation constants are measured in 10% and 20% methanol-water mixtures at 25°C.

For the first dissociation constant the cell used was : -

containing an oxalic acid-sodium hydrogen oxalate buffer mixture. In this cell one electrode is reversible to the hydrogen ion, and the other to the chloride ion. Hence the E.M.F. is given by the equation

$$E = E_0 - \frac{RT}{F} \ln f_{\text{H}} f_{\text{Cl}} m_{\text{H}} m_{\text{Cl}} \dots \dots \dots (1)$$

where E is the measured E.M.F. in volts, and E₀ the normal electrode potential of the cell

and the "m's" and "f's" represent the molal concentrations and activity coefficients respectively, of the ionic species denoted by the subscripts. The thermodynamic dissociation constant K₁ of oxalic acid is given by the equation

$$K = \frac{f_{\text{H}} f_{\text{HC}_2\text{O}_4}}{f_{\text{H}_2\text{C}_2\text{O}_4}} \cdot \frac{m_{\text{H}} m_{\text{HC}_2\text{O}_4}}{m_{\text{H}_2\text{C}_2\text{O}_4}} \dots \dots \dots (2)$$

Eliminating m_H from equations (1) and (2) and rearranging : -

$$E - E_0 + \frac{RT}{F} \ln \frac{m_{H_2C_2O_4} m_{Cl}}{m_{H^+} C_2O_4} = - \frac{RT}{F} \ln \frac{f_H f_{Cl} f_{H_2C_2O_4}}{f_H f_{HC_2O_4}} \\ - \frac{RT}{F} \ln K_1 \dots \dots \dots (3)$$

$$\text{or } \frac{(E - E_0)F}{2.3026RT} + \log \frac{m_{H_2C_2O_4} m_{Cl}}{m_{HC_2O_4}} = - \log \frac{f_H f_{Cl} f_{H_2C_2O_4}}{f_H f_{HC_2O_4}} \\ - \log K_1 \dots \dots \dots (4)$$

This equation is easy to use if the required E_0 values are known at the desired temperatures. The first term on the right contains the logarithm of the ratio of two ionic activity coefficient products. This ratio equals unity at infinite dilution and varies little from unity at low ionic strength. Further, the logarithm of such a quantity as this ratio has been shown by Harned and Robinson,³¹ and by Harned and Owen,¹⁸ to vary linearly with the ionic strength. If therefore, the quantity on the left side of the equation be plotted against the ionic strength (μ) of the solution, a straight line should be obtained. The intercept at $\mu = \text{zero}$ will be $-\log K_1$ (or pK_1 by analogy with pH and $-\log H^+$).

The second term on the left-hand side must now be evaluated; $m_{H_2C_2O_4}$ and $m_{HC_2O_4}$ represent the true molalities of undissociated oxalic acid and hydrogen oxalate ion in the solution. If the unknown hydrogen ion concentration is M_H , and m_1 and m_2 are the stoichiometric molalities of oxalic

acid and sodium hydrogen exalate, assuming complete dissociation of the latter salt: -

$$m_{H_2C_2O_4} = m_1 - m_H$$

$$m_{HC_2O_4} = m_2 + m_H$$

Also assuming complete dissociation of the Sodium Chloride

$$m_{Cl} = m_3$$

Thus the molality term in equation (4) becomes:

$$\log \frac{(m_1 - m_H) m_3}{m_2 + m_H}$$

The only unknown is m_H which must be evaluated by arithmetical approximation from the original equation

$$K_1 = \frac{m_H m_{HC_2O_4}}{m_{H_2C_2O_4}} \cdot \frac{f_H f_{HC_2O_4}}{f_{H_2C_2O_4}}$$

which may be written:

$$K_1 = \frac{m_H (m_2 + m_H)}{m_1 - m_H} \cdot \frac{f_H f_{HC_2O_4}}{f_{H_2C_2O_4}} \dots \dots \dots (5)$$

Following the method of Harned and Embree²³ the activity coefficient ratio in this equation is replaced by the value of $f_H f_{Cl}$ for .01 molal HCl in the presence of Sodium Chloride, calculated from data given by Harned³². The values

for .01 molal HCl can be safely used since Harned and Robinson³¹ have shown that, so long as m_H is .01 molal or less, the values of the activity coefficient of hydrochloric acid in salt solutions are practically independent of m_H and depend only on the ionic strength. Harned and Embree consider that this is sufficiently accurate in the case of formic acid, as any error should be eliminated in the final extrapolation to zero ionic strength. This assumption is the weakest point in the method for as strong an acid as oxalic acid, and may cause a constant error in the results, although not of a very large order. The quadratic equation (5) is then solved for m_H , giving the complete left-hand side of equation (4), i.e.,

$$\frac{(E - E_0)F}{2.3026RT} + \log \frac{(m_1 - m_H)m_3}{m_2 + m_H} = - \log K_1'$$

$$= pK_1'$$

These values of pK_1' , as calculated for solutions of varying strength, are plotted against μ , and a good straight line can be extrapolated to zero ionic strength, where pK_1' becomes equal to pK_1 , and a value of K_1 is obtained. This value is resubstituted in equation (5), a new value of m_H obtained, and the process repeated. Such approximations are carried out until the assumed value of K_1 used in equation (5) gives itself back from the extrapolation of the plot of pK_1' . This value is the true thermodynamic

dissociation constant at the temperature employed.

The method for the 2nd dissociation constant is essentially similar, and is based on that used by Nims²¹ to measure the second dissociation constant of phosphoric acid.

The cell used was the same type of concentration cell without liquid junction, i.e.,

whose E.M.F. is

$$E = E_0 - \frac{RT}{F} \ln f_{\text{H}} f_{\text{Cl}} m_{\text{H}} m_{\text{Cl}}$$

Where E_0 is as above.

$$\text{now, } K_2 = \frac{f_{\text{H}} f_{\text{C}_2\text{O}_4}}{f_{\text{HC}_2\text{O}_4}} \cdot \frac{m_{\text{H}} m_{\text{C}_2\text{O}_4}}{m_{\text{HC}_2\text{O}_4}}$$

and eliminating m_{H} as before:

$$\begin{aligned} -\log K_2 &= \frac{(E - E_0)F}{2.3026 RT} + \log \frac{m_{\text{HC}_2\text{O}_4} m_{\text{Cl}}}{m_{\text{C}_2\text{O}_4}} \\ &+ \log \frac{f_{\text{HC}_2\text{O}_4} f_{\text{Cl}}}{f_{\text{C}_2\text{O}_4}} \dots\dots\dots (6) \end{aligned}$$

By applying the Debye-Huckel formula,

$$-\log f_{\pm}^* = z_+ z_- A \sqrt{\mu}$$

where z_+z_- are the valences of the ions and A and B are constants for the solvent, and ~~solvent~~ and solute respectively, which approximation is accurate enough, seeing that extrapolation is carried out to infinite dilution, the last term in equation (6) is given by : -

$$\log \frac{f_{Cl} f_{HC_2O_4}}{f_{C_2O_4}} = 2A \sqrt{\mu}$$

This method could not be applied to the first dissociation constant, since the value of μ still depends on the unknown m_H which is large for the first dissociation. In this case all the substances present are salts, and considered fully dissociated, and K_2 so small that the concentration of hydrogen ions can be neglected compared with the salt molalities. Hence (6) becomes : -

$$pK_2 = \frac{(E - E_0)F}{2.303RT} + \log \frac{m_1 m_3}{m_2} + 2A \sqrt{\mu}$$

and can readily be calculated. A is the theoretical constant of the Debye-Huckel equation, Hence on plotting the right-hand side of equation (7) against ionic strength, the values of pK_2' obtained should theoretically, and do in practice, lie on a straight line, the intercept of which at zero ionic strength gives pK_2 , and hence a true value of K_2 is obtained. It is important to note that any error in using the Debye-Huckel equation becomes progressively less as the concentration decreases.

Since Hydrogen and chloride ions are present in the cells used, it is necessary to be sure that any complications due to association of hydrochloric acid ions are avoided. The mean distance of approach of ions of hydrogen chloride has been shown to remain constant in these mixtures by Harned and Thomas⁶, the value in water as determined by Harned and Ehlers⁷ being $4.3 \text{ \AA}$. According to Bjerrum's theory of ionic association, if $a > q$ the electrolyte will be completely dissociated. $q = 3.5 \text{ \AA}$ in water and is $< 4 \text{ \AA}$ in 20% methanol-water mixture (Harned and Thomas⁸), hence no association occurs and the Debye-Huckel equation may safely be used for the purpose of extrapolation.

EXPERIMENTAL METHOD

Materials.

Oxalic Acid was purified by two recrystallizations of Hopkins and Williams "Analar" from distilled water. The crystals were drained on a suction pump and dried in a desiccator.

Sodium Hydrogen Oxalate: Previous work³³ in this laboratory has shown, from a phase rule study of the system $\text{H}_2\text{C}_2\text{O}_4 \cdot 2\text{H}_2\text{O} - \text{Na}_2\text{C}_2\text{O}_4 - \text{H}_2\text{O}$, that Sodium Hydrogen Oxalate crystallizes out pure (as $\text{NaHC}_2\text{O}_4 \cdot \text{H}_2\text{O}$) from solutions containing oxalic acid and sodium oxalate in very approximately equimolecular quantities, and that the crystals may be washed without decomposition. Hence the salt was crystallized from a solution equimolecular in A. R. sodium oxalate and A. R. oxalic acid, the crystals drained on a suction pump and washed. These crystals were then recrystallized from distilled water, and dried in a desiccator until their weight remained constant. The sample obtained in this way was found on analysis with a standard potassium permanganate solution to be $100 \pm 0.1\%$ pure.

Sodium Oxalate: The same work³³ has shown that pure crystals will not be obtained from an acid solution of the salt. Since the pure salt solution is alkaline by hydrolysis, so long as too much washing is not carried out, a pure sample is obtained. B.D.H.A.R. sodium oxalate was recrystallized twice from

distilled water. The crystals were dried in an air oven for several hours at 120°C , (as recommended in Cumming and Kays' quantitative analysis) until their weight remained constant.

Sodium Chloride was purified by two recrystallizations of Hopkin and Williams' 'Analar' from distilled water. These crystals were also dried in an air oven.

Water: The solutions were made up with distilled water.

Methyl Alcohol: Kahlbaum's 'Pure' methyl alcohol was purified by fractional distillation with an efficient still head 50 cm. high. The middle fraction of the distillate was refluxed for several hours over a water-bath with 1% by weight of calcium turnings. This was then redistilled. The liquid distilled over at a constant temperature, the first and last 100 cc being discarded. This methyl alcohol had a B.P. of 64.6°C , and a refractive index, measured by a calibrated Abbé refractometer of 1.3284 at 20°C . The value given by Weisberger and Proskauer³⁴ is 1.3286, so the impurity was considered to be water in quantity negligible for consideration in making up solutions. Stock solutions were made up 10% CH_3OH and 20% CH_3OH by weight in approximately five litre lots, the liquids being weighed on a balance accurate to .01 gram. These solutions were kept in air-tight bottles, and were drawn off as required by a siphon, air being admitted through a calcium chloride drying tube, to ensure that the

proportions were not altered by absorption of water from the air.

Hydrogen: Pure electrolytic hydrogen was obtained from a cylinder. It was forced through the cells by means of an aspirator, so that it was already saturated with water vapour at room temperature. The hydrogen was passed over red-hot platinized asbestos to remove any oxygen, and then through a saturator containing the cell solution immersed in the thermostat.

Solutions: Solutions were made up as follows:

The weights used were calibrated by the usual methods against the laboratory standard, and, on the balance used, could be considered accurate to 0.0001 gm. The solutions were made up approximately equal in the molalities of the components, as a 1:1 buffer mixture is ideal experimentally. For the first dissociation constant, equimolecular weights of oxalic acid and sodium hydrogen oxalate were accurately weighed into a weighed quantity of water. This solution was analysed for total oxalate with standard potassium permanganate, and found to agree within 0.1% of the calculated value. To this solution was then added an equimolecular quantity of sodium chloride. In calculating the molalities, allowance was made for the water of crystallization, $\text{H}_2\text{C}_2\text{O}_4 \cdot 2\text{H}_2\text{O}$, and $\text{NaHC}_2\text{O}_4 \cdot \text{H}_2\text{O}$. For the second dissociation constant, equimolecular weights of sodium hydrogen oxalate, sodium

oxalate and sodium chloride were used. The solutions in the alcohol mixtures were made up in exactly the same manner, but were not analysed with potassium permanganate. The strongest solutions used were .05 molal in each salt, except in the case of the second dissociation constant in 20% methanol solution, when the saturated solution was slightly weaker.

The solutions used in the cells were made by diluting weighed quantities of the above stock solutions with weighed quantities of the solvent, water, 10% methanol, or 20% methanol respectively.

Silver Chloride Electrodes.

These were of the 'Jahn' type as used by MacInnes and Beattie³⁵, Allmand and Hunter³⁶, and others. They consisted of .75 sq. cm. squares of platinum gauze welded to platinum wire which in turn made contact with mercury through glass tubes. These were plated with silver by deposition from cyanide solution (7 gms. AgNO_3 + 10 gms. KCN in 200 cc distilled water), 2 milliamps per electrode being passed for 16 hours. They were washed in running tap water and further washed in several portions of boiling distilled water. They were then chloridized anodically in .1N HCl solution for half an hour, with a current density of 6 milliamps per electrode, as recommended by Allmand and Hunter³⁶. Electrodes so prepared agreed to .1 mV. except on a few occasions, generally

due to cracks appearing in the seal. Such electrodes were cleaned as follows : - Silver chloride was removed by .880 ammonia solution, then the silver by boiling with concentrated nitric acid. The final cleansing was after the method of Morgan, Lammert, and Campbell³⁷, by immersion in a cold concentrated sulphuric acid - dichromate cleaning mixture, prepared from C.P. chemicals, which was then heated to 125°C, and allowed to cool to room temperature. The electrodes were then rinsed with tap water and left standing in distilled water until required for plating. Electrodes were used once only to avoid any possible poisoning.

Hydrogen Electrodes.

Platinum wires were sealed into 4 mm tubes, and mounted in pairs in rubber stoppers. Old deposits of platinum black were removed with a damp cloth, followed by polishing with jeweller's rouge. The electrodes were then cleaned in chromic acid in the same way as the silver chloride electrodes. The platinum black was deposited from a 2% solution of platinum chloride, containing .02 gm. lead acetate per 100 c.c; .3 ampere per sq. cm. of cathode surface being passed for one minute as recommended by Harned³⁸. The electrodes were then hydrogenated as cathodes in a 7% sulphuric acid solution for half an hour or longer to remove occluded chlorine, rinsed thoroughly in tap water, and kept in distilled water till required. The electrodes were used once only then cleaned

FIG. I

and replatinized. All electrodes were rinsed with the cell solution before use.

Cells.

The cells used were of a very simple type (as shown in Fig. I). The stopper A carried a glass tube B, which contained two hydrogen electrodes CC, and two silver-silver chloride electrodes DD. The hydrogen entered through E and escaped through holes FF about 1 cm. from the bottom of the tube. The cell was made airtight with collodion, except for an outlet through a capillary tube G about 40 cm. long, which allowed the hydrogen to escape, and prevented any back diffusion of oxygen. This device is considered quite satisfactory by Clark³⁹. The cells were filled to about 1 cm. below the bottom of the stopper, and then immersed in the thermostat to the level of the stopper. Hydrogen was admitted to the electrode tube at the rate of about eight pulsations every five seconds (Clark³⁹), so that the electrodes were alternately bathed in solution and exposed to the hydrogen. Unless the rate was very slow, the potentials were independent of the rate of bubbling of the hydrogen. Harned and his co-workers keep the hydrogen and silver-silver chloride electrodes in separate compartments to prevent the hydrogen ^a affecting the silver-silver chloride electrode. However, previous work in this laboratory has shown that this effect cannot be measured on the potentiometer used. Hence, the present cell was used

which had the great advantages of ease of manipulation, and a low resistance, which allowed the E.M.F.'s to be measured accurately down to the most dilute solutions.

When the hydrogen was turned on, the potential between the two hydrogen electrodes was measured. This fluctuated for about 20 minutes, until equilibrium was obtained, when the pair invariably gave a zero potential difference. The pair of silver-silver chloride electrodes were not quite so consistent, but the electrodes were discarded if a potential difference greater than $\cdot 1$ mv. developed. The cells generally reached equilibrium in an hour when all dissolved oxygen had been removed by the hydrogen bubbles (which at the same time kept the solution stirred). Combining the two hydrogen and two silver-silver chloride electrodes, four readings could be taken and the mean of these readings measured quarter hourly was taken. The cells remained constant for about three hours after equilibrium was obtained, and then started to drift downwards. For the initial measurements duplicate cells were used, giving eight readings in all, but since these agreed to within $\cdot 1$ mv., the limit of the apparatus error, the later e.m.f.'s were measured in one cell only.

Between runs the cells and hydrogen tubes were cleaned in a chromic acid bath, in the same manner as the electrodes, washed with tap water, finally with distilled water, and dried in an air-oven.

During each reading, the barometric pressure was observed, and the E.M.F.'s were corrected to 1 atmosphere pressure of pure hydrogen by means of data given by Clark³⁹. Similar corrections were calculated for the methanol-water mixtures, taking the vapour pressures of 10% CH₃OH - H₂O, and 20% CH₃OH - H₂O mixtures as 33.7 mm. and 43.8 mm. of mercury respectively, as measured by Harned and Thomas⁶. This correction was of the order of about .2 - .6 m. and it was considered sufficiently small to use the vapour pressures of the pure solvents, and to neglect any back pressure along the capillary tube, or the 2 - 3 cm. head of solution under which the hydrogen bubbled out. A correction was also applied to the E.M.F.'s for the change in E.M.F. of the Weston cadmium cell with temperature, the correction being slightly less in value than that for the pressure correction.

The first dissociation constant in water was measured at 25°C, 30°C, and 35°C, the first and second dissociation constants in the alcohol-water mixtures at 25°C only. The water solutions were reproducible to .1 m. but the alcohol solutions varied more, and were only reproducible to .2 m.

Other Apparatus.

A gas heated thermostat was used, controlled by a mercury-chloroform regulator to the required temperatures, 25°C ± .005, 30°C ± .005, 35°C ± .01, recorded by a thermometer

calibrated against the laboratory standard. The standard thermometer could only be read to $\cdot 05^{\circ}$, but the fluctuations in the thermostat were recorded by a Beckmann thermometer.

A Cambridge Instrument Co. potentiometer was used with a range of 0 - 1.8 volts. It was accurate to $\cdot 05$ mv. according to two Weston normal cells made by the Cambridge Instrument Co., one of which was used as a standard.

A Cambridge Instrument Co. galvanometer of the D'Arsonval type was used with a sensitivity of 500 mm. scale divisions per micro-ampere.

CALCULATION OF RESULTS

First Dissociation Constant

The formula used in calculating the First Dissociation constant is : -

$$\frac{(E - E_0)F}{2.3026RT} + \log \frac{m_{H_2C_2O_4} m_{Cl}}{m_{HC_2O_4}}$$

$$= - \log \frac{f_H f_{Cl} f_{H_2C_2O_4}}{f_H f_{HC_2O_4}} - \log K_1 = pK_1 \dots\dots\dots (A)$$

The first term on the left contains E_0 , which is known, and E which is measured. In the second term, the true molalities of the undissociated oxalic acid, $m_{H_2C_2O_4}$, and of the hydrogen oxalate ion, $m_{HC_2O_4}$, must be evaluated by arithmetical approximation. This was found necessary for acetic¹⁹, formic²³, and chloracetic²⁶ acids, and since oxalic acid in its first dissociation is considerably stronger than any of these acids, the same approximation must be carefully carried out. The first term on the right-hand side was found to vary linearly with ionic strength, hence by plotting the left-hand side against the ionic strength, the line can be readily extrapolated to zero ionic strength, where the intercept gives $-\log K_1$, or pK_1 .

The method of approximation as used for acetic acid

was employed here to obtain the true molalities of the undissociated acid, hydrogen oxalate and chloride ions from the stoichiometrical molalities m_1 , m_2 , m_3 , of oxalic acid, sodium hydrogen oxalate, and sodium chloride. The sodium chloride is assumed completely dissociated hence $m_{Cl} = m_3$.

In the first approximation the value obtained in this laboratory last year, $K_1 = .01718$, was used in the equation:

$$K_1 = \frac{(m_2 + m_H) m_H}{m_1 - m_H} f_H f_{Cl} \dots\dots\dots (B)$$

The values of $f_H f_{Cl}$ in sodium chloride solutions were calculated from data given by Harned.³²

His figures are for the cell

These are combined with the cell

for which Harned and Ehlers⁷ give

$$E_{25^\circ C} = + .46416 \quad \text{and} \quad E_o = .22239$$

$$\text{Hence } .46416 = .22239 - \frac{2RT}{F} \ln f_{HCl} \times .01$$

$$\text{and } \therefore f_{HCl} = .904$$

Now for Cell (1), $E_m = E_o - .05915 \log f_H \times .01 \times f_{Cl} (.01 + m)$

and for Cell (2) $.46416 = E_o - .05915 \log .904^2 \times .01^2$

subtracting

$$.46416 - E_m = .05915 \log \frac{f_H f_{Cl} \times .01 \times (.01 + m)}{.904^2 \times .01^2}$$

which gives $f_H f_{Cl}$ and $f_{HCl} = \sqrt{f_H f_{Cl}}$

The logarithms of the values of f_{HCl} when plotted against m gave a reasonably smooth curve from which f_{HCl} was read off at the required concentrations, and the square of this value gave $f_H f_{Cl}$.

From further data of Harned³² calculated from the formula given by him for the cell (1) with change of temperature, of the type

$$E = E_{25^{\circ}C} + a(t - 25) + b(t - 25)^2$$

the corresponding activity coefficient product was calculated at 30°C and 35°C.

No data were available for such cells in methanol-water mixtures, the only figures being for the .01M acid with no sodium chloride present. The values at 25°C as given by Harned and Thomas⁸ being:

f_{HCl}	pure H_2O	=	•904
f_{HCl}	10% CH_3OH	=	•897
f_{HCl}	20% CH_3OH	=	•898

It was considered sufficient to reduce the values used for the activities in pure water in the same proportions as the above to give those in the methanol-water mixtures. The error in taking these values is certainly very small, and should have no effect in the extrapolation to infinite dilution.

The quadratic equation (B) is then solved for m_{H} which gives the true molalities: -

$$m_{\text{H}_2\text{C}_2\text{O}_4} = m_1 - m_{\text{H}}$$

$$m_{\text{H}_2\text{C}_2\text{O}_4} = m_2 + m_{\text{H}}$$

Substituting these values in the left-hand side of (A) gives a value of pK_1' , and the plot of these points extrapolated gives a new value of pK . This value of K is then used in B to give a better approximation to m_{H} , and the process repeated, until a chosen value of K gives back itself within the limits of the graphical extrapolation.

The ionic strength was calculated as follows from the Lewis and Randall formula

$$\mu = \frac{1}{2} \sum m z^2$$

where m = molality of each ion, and z the valence and the sum being taken for all the ions present. In this case, there are present oxalic acid m_1 , which has dissociated to give m_H hydrogen ions, sodium hydrogen oxalate, m_2 , and sodium chloride, m_3 , both as salts considered completely dissociated.

$$\begin{array}{llll} \text{i.e., the ions are} & \text{H}^+ & = & m_H \\ & \text{HC}_2\text{O}_4^- & = & m_2 + m_H \\ & \text{Na}^+ & = & m_2 + m_3 \\ & \text{Cl}^- & = & m_3 \end{array}$$

$$\begin{aligned} \therefore \mu &= \frac{1}{2} m_H (1)^2 + \frac{1}{2} (m_2 + m_H) (1)^2 + \frac{1}{2} (m_2 + m_3) (1)^2 \\ &\quad + \frac{1}{2} m_3 (1)^2 \\ &= m_2 + m_3 + m_H \end{aligned}$$

Great care is needed in carrying out these approximations. When the value of K is distant from the true values, the points are very scattered around the straight line, especially in the dilute solutions where a change of 0.0001 in the molality of the undissociated acid is of the order of 2% of the concentration, and causes large changes in pK . Hence, owing to the difficulty of placing the

straight line, a fortuitous agreement may be obtained between two successive approximations which leads to the approximations not being carried far enough.

This is shown in the value obtained in this laboratory last year, $K_1 = .01718$, which is recalculated in the present work. The approximations can be continued to change the value to $K_1 = .02884$. A further error was discovered in the activities used, making the final recalculated value $K_1 = .05023$. The recalculation and the original values are shown in Graph I, and the recalculation can be seen to be much more accurate. The plotting of the line could not be carried out with an accuracy of more than $\pm .002$ in pK , which is probably about the same order as the experimental error.

As more skill was gained in using the approximation method, time could be saved by estimating values of K , and by calculating only three or four points when the approximate value was obviously remote from the true value. Thus, for the value at 25°C , twelve approximations were needed, at 30°C seven approximations, at 35°C , four approximations, in 10% CH_3OH , six approximations, in 20%, five approximations, and for the recalculation, four approximations.

Constants used in the calculations.

$$\frac{2.3026RT}{F} = .05915 \quad \text{at} \quad 25^\circ\text{C}$$

$$\frac{2 \cdot 3026RT}{F} = .06014 \text{ at } 30^{\circ}\text{C}$$

$$\frac{2 \cdot 3026RT}{F} = .06113 \text{ at } 35^{\circ}\text{C}$$

For the cell

In pure water, Harned & Ehlers⁷ give:

$$25^{\circ}\text{C} \quad E_0 = + .2224$$

$$30^{\circ}\text{C} \quad E_0 = + .2191$$

$$35^{\circ}\text{C} \quad E_0 = + .2156$$

In methanol-water mixtures at 25°C , Harned & Thomas⁶ :

$$10\% \text{ CH}_3\text{OH} - 90\% \text{ H}_2\text{O} \quad E_0 = + .2154$$

$$20\% \text{ CH}_3\text{OH} - 80\% \text{ H}_2\text{O} \quad E_0 = + .2088$$

For the cell $\text{Pt} \mid \text{quinhydrone} \mid \text{HCl} \mid \text{AgCl} \mid \text{Ag}$
(used for recalculation)

$$25^{\circ}\text{C} \quad E_0 = - .4774 \quad (\text{Harned and Wright}^{40})$$

The values obtained for the first dissociation constant were: -

25°C	pK	=	1.336	K	=	.04613	) water
25°C	pK	=	1.299	K	=	.05023 (recalc)	
30°C	pK	=	1.258	K	=	.05521	
35°C	pK	=	1.271	K	=	.05358	
25°C	pK	=	1.514	K	=	.03062	10% $\text{CH}_3\text{OH} - \text{H}_2\text{O}$
25°C	pK	=	1.647	K	=	.02254	20% $\text{CH}_3\text{OH} - \text{H}_2\text{O}$

Second Dissociation Constant.

The formula in this case is: -

$$- \log K_2 = \frac{(E-E_0)F}{2 \cdot 3026 RT} + \log \frac{m_{\text{HC}_2\text{O}_4} m_{\text{Cl}}}{m_{\text{C}_2\text{O}_4}} + 2A \sqrt{\mu}$$

The right-hand side of this equation is plotted against the ionic strength, and the straight line so obtained extrapolated to zero ionic strength; the intercept is $-\log K_2$ or $\text{p}K_2$.

The calculation is as before, but no approximations are needed.

The molalities can be taken as equal to the stoichiometrical molalities of sodium hydrogen oxalate, sodium oxalate, and sodium chloride, since, as salts of a strong base, they can be considered completely dissociated. Also the second dissociation of the hydrogen oxalate ion can be considered as infinitely small compared with the concentrations employed.

Calculating the ionic strength we have:

$$\text{i.e. } \text{Na}^+ = m_1 + 2m_2 + m_3$$

$$\text{HC}_2\text{O}_4^- = m_1$$

$$\text{C}_2\text{O}_4^{2-} = m_2$$

$$\text{Cl}^- = m_3$$

$$\begin{aligned}
 \therefore \mu &= \frac{1}{2} (m_1 + 2m_2 + m_3) (1)^2 + \frac{1}{2} m_1 (1)^2 + \frac{1}{2} m_2 (2)^2 \\
 &\quad + \frac{1}{2} m_3 (1)^2 \\
 &= m_1 + 3m_2 + m_3
 \end{aligned}$$

The values of A at 25°C used was $.506$ as given by Clark³⁹.

In CH_3OH mixtures, Harned and Thomas⁸ have shown $A = K'a$, where a the mean distance of approach of the ions is the same as in water, and $K' \propto \frac{1}{D K}$, where D is the dielectric constant.

Using the values:

D pure H_2O	=	78.54	Wyman, ⁴¹
D 10% CH_3OH	=	74.05	Akerlof ⁴²
D 20% CH_3OH	=	69.20	Akerlof ⁴²

Hence	A	10% CH_3OH	=	.521	552
	A	20% CH_3OH	=	.539	611

These values of pK_2 gave an excellent straight line, the extrapolated values being at 25°C : -

10% CH_3OH	-	90% H_2O	$pK_2 =$	4.553	$K =$	2.779
				4.538		2.897×10^{-5}
20% CH_3OH	-	80% H_2O	$pK_2 =$	4.822	$K =$	1.417
				4.808		1.556×10^{-5}

Tables.

Tables I, II, III, IV, V, record the values for the first dissociation constant under varying conditions. The first three columns give the molalities m_1 for $\text{H}_2\text{C}_2\text{O}_4$, m_2 for

NaHC_2O_4 , m_3 for NaCl . The fourth column gives the calculated activity product $f_{\text{H}} f_{\text{Cl}}$ in the salt solution, the fifth, the calculated hydrogen ion molality, the sixth the calculated ionic strength. The seventh gives the electromotive force in volts corrected for temperature and pressure variations, and the eighth column the calculated values of

$$\frac{(E - E_0)F}{2 \cdot 3026RT} + \log \frac{m_{\text{H}_2\text{C}_2\text{O}_4} m_{\text{Cl}}}{m_{\text{HC}_2\text{O}_4}} = \text{pK}_1'$$

These results are those for the final approximations. Table VI gives the recalculated values from the results obtained last year, in this laboratory. Unfortunately, the E.M.F.'s were not corrected for change in E.M.F. of the standard cell with temperature, so this result cannot be regarded as strictly accurate.

Tables VII and VIII are for the second dissociation constant, the figures being in this case m_1 for NaHC_2O_4 , m_2 for $\text{Na}_2\text{C}_2\text{O}_4$, and m_3 for NaCl , and the final column

$$\text{pK}_2' = \frac{(E - E_0)F}{2 \cdot 3026RT} + \log \frac{m_{\text{HC}_2\text{O}_4} m_{\text{Cl}}}{m_{\text{C}_2\text{O}_4}} + 2A\sqrt{\mu}$$

Graphs.

Graph I shows the plots for the first dissociation constant at varying temperatures; A, B, C being for 25°C , 30°C , and 35°C respectively. E represents the value obtained

in this laboratory last year and D the recalculation of those results. D and C should be between A and B, but are shown in their present positions for clarity. Graph II shows the plots for the first dissociation constant at 25°C in pure water, 10% methanol-water, and 20% methanol-water, H, G, F, respectively. Graph III shows the similar plots for the second dissociation constant, K, J, I. K is the value obtained last year by the previous worker. Graph IV shows on the left-hand side the plot of K_1 against temperature; on the right-hand side the plots of pK_1 and pK_2 against the reciprocals of the dielectric constants of the various solvents. The black circles are the results obtained last year.

TABLE I

1st Dissociation Constant in H₂O

Temp. = 25°C

Assuming $K_1 = .04571$ or $pK_1 = 1.340$

m_1	m_2	m_3	$f_H f_{Cl}$	m_H	μ	E corr.	pK_1
.04986	.04999	.05097	.667	.02398	.12494	+ .4049	1.33643
.03293	.03302	.03366	.696	.01846	.08514	+ .4214	1.34028
.02484	.02491	.02540	.714	.01526	.06557	+ .4323	1.35091
.01661	.01648	.01704	.738	.01145	.04497	+ .4498	1.34235
.01263	.01266	.01291	.752	.00928	.03485	+ .4613	1.33362
.00828	.00822	.00850	.769	.00662	.02334	+ .4798	1.33294
.00643	.00645	.00657	.780	.00535	.01837	+ .4914	1.32689
.00404	.00402	.00415	.790	.00357	.01174	+ .5122	1.30914
.00197	.00195	.00202	.803	.00184	.00581	+ .5469	1.32674
Extrapolated					0		1.336

TABLE II

1st Dissociation Constant in H₂O

Temp. = 30°C

Assuming $K_1 = .05495$ or $pK_1 = 1.260$

m_1	m_2	m_3	$f_H f_{Cl}$	m_H	μ	E corr.	pK_1
.04986	.04999	.05097	.666	.02596	.12692	+ .4046	1.28967
.03293	.03302	.03366	.694	.01976	.08644	+ .4211	1.28306
.02484	.02491	.02540	.712	.01621	.06652	+ .4325	1.27518
.01661	.01648	.01704	.736	.01202	.04554	+ .4497	1.27283
.01263	.01266	.01291	.750	.00968	.03525	+ .4615	1.26228
.00828	.00822	.00850	.767	.00684	.02356	+ .4800	1.24818
.00620	.00626	.00630	.778	.00533	.01789	+ .4938	1.24247
.00332	.00329	.00340	.794	.00303	.00972	+ .5234	1.25304
.00197	.00195	.00202	.801	.00186	.00583	+ .5492	1.25470
Extrapolated					0		1.258

TABLE III1st Dissociation Constant in H_2O Temp. = $35^{\circ}C$ Assuming $K_1 = .05309$ or $pK_1 = 1.275$

m_1	m_2	m_3	$f_H f_{Cl}$	m_H	μ	E. corr.	pK_1
*04973	*05025	*05060	*661	*02557	*12642	+ *4045	1.29760
*03321	*03356	*03379	*689	*01964	*08699	+ *4203	1.28408
*02476	*02502	*02519	*707	*01600	*06621	+ *4332	1.29037
*01660	*01677	*01689	*733	*01189	*04555	+ *4505	1.28602
*01236	*01249	*01258	*752	*00943	*03450	+ *4632	1.27609
*00828	*00837	*00842	*767	*00679	*02358	+ *4814	1.26591
*00620	*00626	*00630	*776	*00530	*01786	+ *4951	1.26285
*00414	*00418	*00421	*788	*00370	*01210	+ *5142	1.25588
*00332	*00329	*00340	*794	*00303	*00972	+ *5249	1.25287
Extrapolated					0		1.271

TABLE IV1st Dissociation Constant in 10% CH_3OH - 90% H_2O Temp. = $25^{\circ}C$ Assuming $K_1 = .03090$ or $pK_1 = 1.510$

m_1	m_2	m_3	$f_H f_{Cl}$	m_H	μ	E. corr.	pK_1
*05036	*05099	*05151	*656	*02007	*12257	+ *4013	1.47727
*03224	*03264	*03297	*684	*01558	*08119	+ *4192	1.50205
*02509	*02541	*02567	*702	*01334	*06442	+ *4291	1.49726
*01597	*01617	*01633	*728	*00988	*04238	+ *4475	1.50572
*01256	*01272	*01285	*746	*00832	*03389	+ *4582	1.51804
*00801	*00811	*00819	*762	*00594	*02224	+ *4769	1.50254
*00628	*00636	*00642	*771	*00490	*01764	+ *4872	1.49097
*00407	*00412	*00416	*781	*00343	*01171	+ *5063	1.46900
*00311	*00315	*00318	*785	*00270	*00904	+ *5183	1.46894
Extrapolated					0		1.514

TABLE V1st Dissociation Constant in 20% CH₃OH - 80% H₂O

Temp. = 25°C

Assuming $K_1 = .02239$ or $pK_1 = 1.650$

m_1	m_2	m_3	f_H	f_{Cl}	m_H	μ	E. corr.	pK_1'
.05196	.05288	.04927	.643	.01724	.11939	+	.4007	1.63161
.03641	.03705	.03452	.671	.01433	.08590	+	.4138	1.63705
.02605	.02651	.02483	.689	.01193	.06327	+	.4266	1.64219
.01818	.01850	.01724	.714	.00959	.04533	+	.4403	1.63576
.01297	.01319	.01236	.733	.00769	.03324	+	.4544	1.64708
.00916	.00932	.00869	.748	.00605	.02406	+	.4687	1.63902
.00653	.00664	.00622	.757	.00471	.01757	+	.4834	1.64130
.00477	.00486	.00453	.767	.00366	.01305	+	.4969	1.64165
Extrapolated						0		1.647

TABLE VI1st Dissociated Constant in H₂O

Temp. = 25°C

Assuming $K_1 = .05000$ or $pK_1 = 1.300$

$m_1 = m_2 = m_3$	$f_H f_{Cl}$	m_H	μ	Euncorr.	pK_1'
*C4959	*664	*03492	*12410	- *2944	1*30918
*03306	*694	*01916	*08528	- *2784	1*30880
*02480	*711	*01573	*06533	- *2664	1*31148
*01984	*724	*01340	*05308	- *2575	1*30244
*01653	*736	*01167	*04473	- *2500	1*29912
*00992	*762	*00781	*02765	- *2274	1*29863
*00496	*789	*00453	*01425	- *1963	1*27913
*00248	*799	00250	*00726	- *1643	1*26362
Extrapolated			0		1*299

TABLE VII2nd Dissociation Constant in 10% CH₃OH - 90% H₂O. Temp. = 25°C.

m ₁	m ₂	m ₃	μ	E. corr.	pK ₂ '
*04918	*04978	*04910	*24762	+ *5413	4.71405 4.74491
*03325	*03366	*03320	*16743	+ *5536	4.65986 4.68523
*02463	*02493	*02459	*12401	+ *5631	4.63090 4.65277
*01662	*01682	*01659	*08367	+ *5758	4.60904 4.62677
*01234	*01248	*01232	*06210	+ *5847	4.58848 4.60393
*00829	*00839	*00828	*04174	+ *5964	4.56695 4.57762
*00616	*00623	*00615	*03100	+ *6050	4.55407 4.56479
*00415	*00420	*00415	*02090	+ *6174	4.55977 4.56673
*00307	*00310	*00306	*01543	+ *6262	4.55599 4.56364
Extrapolated			0		4.538 4.553

TABLE VIII2nd Dissociation Constant in 20% CH₃OH - 80% H₂O. Temp. = 25°C.

m ₁	m ₂	m ₃	μ	E. corr.	pK ₂ '
*03786	*04026	*03941	*19805	+ *5576	4.94552 5.05441
*02626	*02792	*02733	*13735	+ *5692	4.90252 4.97294
*01862	*01980	*01939	*09741	+ *5803	4.87798 4.95414
*01277	*01358	*01329	*06686	+ *5918	4.85062 4.91371
*00925	*00983	*00963	*04837	+ *6023	4.84688 4.90555
*00657	*00699	*00684	*03438	+ *6116	4.81783 4.86307
*00453	*00481	*00472	*02368	+ *6233	4.82139 4.85974
*00325	*00346	*00339	*01702	+ *6334	4.82201 4.83447
Extrapolated			0		4.808 4.848

GRAPH 1

GRAPH II.

DISCUSSION

For the first dissociation constant, the values in the literature showing the best agreement at 25°C, are .040, Harnel and Butler⁴⁹, by catalysis, .057, Drucker¹¹, by conductivity, .057, Dawson and Smith¹⁰, by catalysis, .059, Gane and Ingold, by electrometric titration, but a wide range from .020 to .17 is obtained by the same types of methods. The present value at 25°C is .046, which shows reasonable agreement with these results, and better agreement with the value .050, recalculated in the present work, but independently obtained in this laboratory. This latter value was obtained by the same method as in the present work, but using a different technique, a quinhydrone electrode, and potassium salts in the cells. The present value .046 is considered more accurate than the value .050, owing to temperature corrections not being applied in the latter case, and since the high resistance cells used previously did not give such accurate E.M.F.'s. It is to be expected that the 'thermodynamic K' would be lower than the 'classical' values obtained by conductivity and catalysis measurements. The Gane and Ingold value is a 'thermodynamic' K but, as pointed out above, the liquid junction errors in the electrometric method may be very large.

However, the present method for the first dissoc-

dissociation constant must be considered very near its limit of applicability for an acid as strong as oxalic. For acetic acid, Harned and Ehlers¹⁹ found that an approximation of $K = 1.65 \times 10^{-5}$ — 1.75×10^{-5} , when used to calculate m_H , was sufficiently accurate to give the true value of K after extrapolation, and Harned and Embree²³ always found three approximations sufficient in the case of formic acid. For oxalic acid, the approximations were very slow, and the method was very sensitive to changes in m_H . The assumption made in the approximation, that $\frac{f_H f_{H_2C_2O_4}}{f_{H_2C_2O_4}}$ can be replaced by $f_H f_{Cl}$ values in similar solutions, may not be justified for a fairly strong acid such as oxalic.

For the second dissociation constant, which is of the same order as those of other acids measured accurately by this method, this difficulty does not arise. The value obtained in this laboratory in pure water 5.012×10^{-5} agrees well with values in the literature. No values in methanol-water mixtures are available for comparison, but, since the method has given such consistent results in similar cases, the values obtained are considered as accurate thermodynamic constants, and, since the method used was the same, can be compared with the value, 5.012×10^{-5} obtained in pure water.

As far as the change in dissociation constant with temperature is concerned, similar results are obtained to

those surveyed by Harned and Embree⁴⁷ for several acids. The variation is not very large, and a maximum is probably obtained at about 32°C, as shown in Graph IV, but not enough temperatures were available to study this effect thoroughly.

When the values of K_1 and K_2 were compared, results were obtained varying little from those calculated using previous values.

Substituting in Bjerrum's equation:

$$\log \frac{K_1}{K_2} - \log 4 = \frac{3.1 \times 10^{-8}}{\gamma}$$

the value obtained is $\gamma = 1.51\overset{0}{A}$, which as pointed out before, is much too low, due principally to neglect of the 'inductive' effect, possibly also of other effects.

The extended equations of Gane and Ingold¹⁴ with the symbols defined as above are:

$$\log \frac{K_1}{K_2} - \log 4 = \frac{q \phi}{2.3026kT}$$

and substituting the values of K_1 and K_2 in this equation

$$\phi = 4.656 \times 10^{-4}$$

γ can only be found now by solving the fifth degree equation

$$\phi = \frac{6.112 \times 10^{-12}}{\gamma} + \frac{1.368 \times 10^{-41}}{\gamma^5}$$

Using Horner's method of solution of such equations,

GRAPH IV

the value of γ correct to one place is 3.5^0 , compared with 3.37^0 quoted by Gane and Ingold¹⁴.

The change in the dissociation constants with change of solvent is as expected in both cases. As the dielectric constant is lowered, the force between the ions is increased, and the dissociation constant falls, as shown by the values : -

	<u>water</u>	<u>10% methanol</u>	<u>20% methanol</u>
K_1	.046	.030	.022
K_2	5.01×10^{-5}	$\frac{2.80}{2.90} \times 10^{-5}$	$\frac{1.42}{1.55} \times 10^{-5}$

Again, as expected, the effect is greater for the second dissociation constant than for the first, because of the doubly charged anion. If the change in dissociation constant is largely dependent on the dielectric constant, as theoretical considerations indicate, the plot of pK against the reciprocal of the dielectric constant should be a straight line. Although three values are not sufficient rigidly to test this relationship, Harned and Embree³⁰, considered this relationship held for acetic acid in these solvents. The plots for oxalic acid are shown in Graph IV. For the first dissociation constant the points deviate slightly from a straight line, but this is to be expected in the case of such a large value for K . The plot of pK_2 gives a good straight line, and hence we may conclude that the

dielectric effect is the predominating one.

The quantitative relationship may therefore be used: -

$$pK_s - pK_w = \frac{Nq^z z}{2.3026RTa} \left(\frac{1}{D_w} - \frac{1}{D_s} \right)$$

$$= \frac{z \times 2.42 \times 10^{-6}}{2a} \left(\frac{1}{D_w} - \frac{1}{D_s} \right)$$

where z is the valence of the anion base, and a is the mean ionic radius, which is calculated.

The values of " a " thus obtained are as follows, with those for acetic acid also, calculated from the figures of Harned and Embree³⁰ from water to: -

	10% methanol	20% methanol
oxalic acid, 1st,	$a = 1.05 \overset{O}{\underset{\underset{A}{ }}{\text{A}}}$	$a = 1.34 \overset{O}{\underset{\underset{A}{ }}{\text{A}}}$
oxalic acid, 2nd,	$a = 1.57 \overset{O}{\underset{\underset{A}{ }}{\text{A}}}$	$a = 1.64 \overset{O}{\underset{\underset{A}{ }}{\text{A}}}$
acetic acid,	$a = 1.46 \overset{O}{\underset{\underset{A}{ }}{\text{A}}}$	$a = 1.32 \overset{O}{\underset{\underset{A}{ }}{\text{A}}}$

The definite differences in " a " as calculated for the change from water to 10% methanol and water to 20% methanol indicate that the assumption that ' a ' is constant in the different solvents is probably unjustified, and that the " a 's" measured are only mean values.

These values may be compared with those obtained by Lynch and La Mer⁵⁰, who used a formula derived in a

similar manner, $2.6\overset{\circ}{\text{A}}$ for acetic, propionic and butyric acids and $1.3\overset{\circ}{\text{A}}$ for benzoic acid. These results were obtained over a wider range of dielectric constant using dioxan-water mixtures. These values are lower than those usually calculated for 'a' from crystal structure measurements, but disagreement is to be expected. In the latter case the inter ionic distances depend solely on Coulomb forces which vary inversally as the dielectric constant. In calculating the values of "a" in solution Coulomb forces only have been taken into account, and Van der Waal's and similar forces between the ions themselves or ions and solvent molecules have been neglected. The values in methanol-water mixtures are still lower, but the results are all of the same order, and the more accurate values for the second dissociation constant for oxalic acid are close to those for acetic acid in the same solvents.

The change of K with dielectric constant has been used by Sorenson, Linderstrom-Lang, Pring and others for analytical purposes. Richardson⁴ used acetone and ethyl alcohol as solvents for the purpose of titrating strong acids in the presence of weak acids. He quotes, without any indication of derivation, the formula:

$$\frac{36000}{T} \left(\frac{1}{D_s} - \frac{1}{D_{H_2O}} \right)$$

for change in pK with change of solvent which is similar to the one used above. His theoretical calculations for ethyl alcohol give the change in pK as 6.8 units for a doubly charged associating ion, and 3.4 for a singly charged ion, i.e., an acid such as acetic should have a pK value 3.4 units higher in ethyl alcohol than in water. The same formula used for methyl alcohol gives the theoretical changes as 4.5 and 2.2 units. If the straight lines of Graph IV are extrapolated to a value of $\frac{1}{D}$ representing pure methyl alcohol, the corresponding changes in pK_2 and pK_1 are 5.5 units and 2.9 units. This shows a reasonable agreement, considering the dubious validity of the extrapolation, and the fact that for the first dissociation the points do not lie very well on a straight line. Harned and Embree³⁰, consider, for acetic acid, that such an extrapolation is not valid beyond 30 in 40% methanol-water mixtures.

Strong acids which are completely dissociated are not affected by change of dielectric constant, hence titration of a mixture of a strong and a weak acid first in water and then in some mixed solvent where the dissociation of the weak acid is repressed, gives a method of analysis for both acids. The results seem to indicate that this method would not be practicable for strong acids in the presence of oxalic acid if methanol was used as solvent. Certainly the second dissociation would be completely repressed, but the

first dissociation would still be appreciable. A solvent of lower dielectric constant such as 80% dioxan-water ($D = 11$) would perhaps be suitable.

The mean heat of ionization of the first dissociation of oxalic acid can be calculated between 25 - 30°C, and between 30 - 35°C by means of the integrated form of the "Isochore".

$$\log \frac{K_2}{K_1} = \frac{\Delta H}{2 \cdot 3026R} \frac{T_2 - T_1}{T_2 T_1}$$

$$\text{or } pK_1 - pK_2 = \frac{\Delta H}{2 \cdot 3026R} \frac{T_2 - T_1}{T_2 T_1}$$

This integration can only be carried out if ΔH is assumed constant over the temperature range employed, hence a mean heat of ionization between the two temperatures is obtained.

For 25 - 30°C

$$1.336 - 1.258 = \frac{\Delta H}{2 \cdot 3026 \times 1.988} \times \frac{5}{298 \times 303}$$

$$\therefore \Delta H = + 6448 \text{ cal.}$$

For 30 - 35°C

$$1.258 - 1.271 = \frac{\Delta H}{2 \cdot 3026 \times 1.988} \times \frac{5}{303 \times 308}$$

$$\therefore \Delta H = - 1111 \text{ cal.}$$

The change in sign follows since K passes through a maximum at approximately 32°C . Sufficient results were not available to calculate ΔC_p , the difference between the specific heats of the undissociated molecule and the ions, from these data, and hence the relationship suggested by Moelwyn-Hughes⁴⁸, that $-\Delta C_p$ is always greater than 30 cal. per degree, could not be tested.

SUMMARY

1. The first dissociation constant of oxalic acid has been determined from electromotive force measurements of cells without liquid junction at 25°, 30° and 35°C.
2. The first and second dissociation constants have been measured by the same method in 10% methanol-water and 30% methanol-water mixtures.
3. These values have been compared, using relationships suggested by Bjerrum, Gane and Ingold, and Bronstead.
4. The application of these results to differential acid titration has been discussed.
5. The mean heat of ionization of the first dissociation of oxalic acid has been calculated between 25° - 30°C, and between 30° - 35°C.

REFERENCES

1. Bronsted, Rec. Trav. Chim. 42, 718. (1923)
2. Lowry, Chem. and Ind. 42, 43. (1923)
3. Bronsted, J. Phys. Chem. 30, 777. (1926)
4. Richardson, Proc. Roy. Soc. B115, 121. (1934)
5. Hartley et al. Chemistry at the Centen. Meeting
Brit. Ass. Adv. Science, (1932)
6. Harned and Thomas, J.A.C.S., 57, 1666, (1935)
7. Harned and Ehlers, J.A.C.S. 55, 2179. (1933)
8. Harned and Thomas, J.A.C.S., 58, 761. (1936)
9. Chandler, J.A.C.S., 30, 713. (1908)
10. Dawson and Smith, J. C. S., 132, 2539. (1929)
11. Drucker, Z. Physikal Chem. 96, 381. (1920)
12. Hamer and Acree, J. Research Natl. Bur. Standards,
17, 605. (1936)
13. Enklaar, Chem. Weekblad, 8, 375, 824. (1911)
14. Cane and Ingold, J. C. S., 134, 2153, (1931)
15. Britton, J. C. S., 127, 1896; (1925)
16. Glasstone, "The Electrochemistry of Sol-
utions, 1st Edition, p. 92.
17. Philbrick, "An Introduction to Physical
Chemistry", p. 187.
18. Harned and Owen, J.A.C.S., 52, 5079. (1930)
19. Harned and Ehlers, J.A.C.S., 54, 1350. (1932)
20. MacInnes and Shedlovsky, J.A.C.S., 54, 1429, (1932)
21. Nims, J.A.C.S., 55, 1946. (1933)
22. Bjerrum & Unmack, Kgl. Danske Videnskab. Selskab.
Math. pp. Medd. 9, 129. (1929)

23. Harned and Embree, J.A.C.S., 56, 1042. (1934)
24. Harned and Ehlers, J.A.C.S., 55, 652. (1933)
25. Harned and Sutherland, J.A.C.S., 56, 2039. (1934)
26. Wright, J.A.C.S., 56, 314, (1934)
27. Nims, J.A.C.S., 58, 987, (1936)
28. Nims, J.A.C.S., 56, 1110. (1934)
29. Hamer, J.A.C.S., 56, 860. (1934)
30. Harned and Embree, J.A.C.S., 57, 1669, (1935)
31. Harned and R.A.Robinson, J.A.C.S., 50, 3157. (1928)
32. Harned, J.A.C.S., 57, 1865. (1935)
33. Cole, Thesis, N.Z.University, 1937.
34. Weissberger and Proskauer, "Organic Solvents". p. 25.
35. MacInnes and Beattie, J.A.C.S., 42, 1117. (1920)
36. Allmand and Hunter, Trans. Farad. Soc. 24, 300. (1928)
37. Morgan Lammert and Campbell, J.A.C.S., 53, 454. (1931)
38. Harned, J.A.C.S., 48, 326. (1926)
39. Clark, "Determination of Hydrogen Ions,"
p. 260.
40. Harned and Wright, J.A.C.S., 55, 4849. (1933)
41. Wyman, Phys. Rev. 35, 623. (1930)
42. Akerlof, J.A.C.S., 54, 4125. (1932)
43. Glasstone, Recent Advances in Phys. Chem.
pp. 447 - 451.
44. Bjerrum, Z. Phys. Chem. 106, 219. (1932)
45. R.A.Robinson, Private Communication.
46. Gurney. "Ions in Solution". p.1.
47. Harned and Embree, J.A.C.S., 56, 1050. (1934)

48. Moelwyn-Hughes, Trans. Farad. Soc. 34, 105,
(1938)
49. Hornel and Butler, J. C.S. 139, 1361. (1936)
50. Lynch and La Mer, J.A.C.S., 60, 1259. (1938)