

**IN SEARCH OF 4 – (12,6,4) DESIGNS:  
PART I**

by

**D.R. Breach**

*Department of Mathematics, University of Canterbury,  
Christchurch, New Zealand.*

and

**Martin J. Sharry and Anne Penfold Street**

*Centre for Combinatorics, Department of Mathematics,  
The University of Queensland, Brisbane 4072, Australia.*

No. 73

January, 1993

To appear in the Australasian Journal of Combinatorics, 7 (1993) pp237–245.

# IN SEARCH OF 4 — (12, 6, 4) DESIGNS: PART I

**D.R.Breach,**

Department of Mathematics, University of Canterbury, Christchurch, New Zealand;

**Martin J. Sharry and Anne Penfold Street,**

Centre for Combinatorics, Department of Mathematics,  
The University of Queensland, Brisbane 4072, Australia.

## **Abstract**

As a first step towards finding all 4-(12, 6, 4) designs which are not 5-(12, 6, 1) designs, it is shown that if such a design has a pair of blocks with five points in common, then there is a unique way of assigning the replicas of the seven points from that pair of blocks to the other blocks of the design.

## **1. Introduction**

A  $t - (v, k, \lambda)$  is a collection of subsets, called *blocks*, of a set  $S$  with  $v$  elements, called *points*, such that every  $t$ -subset of  $S$  is contained in precisely  $\lambda$  blocks. If  $s$  is a whole number such that  $0 \leq s \leq t$ , then a  $t$ -design is also an  $s$ -design with, of course, a different  $\lambda$  value. Thus the 5-(12, 6, 1) design, which is unique in structure, is also a 4-(12, 6, 4) design. However the converse is not true; there are 4-(12, 6, 4) designs which are not 5-(12, 6, 1) designs. Nine non-isomorphic such designs are constructed in Breach, Elmes, Sharry and Street [1]. In this series of papers we show, by a different construction, that, together with the 5-(12, 6, 1) design, these are the only 4-(12, 6, 4) designs. The work proceeds in three major steps. In Part I it is shown that if the design is not a 5-design, then seven of the points determine a unique skeleton for a 4-(12, 6, 4) design. In Part II [2] it is shown that 32 blocks of the skeleton can be completed in a unique fashion, and embedded in these blocks are the 30 blocks of a 3-(10, 4, 1) design. In Part III [3] it is shown that the 100 partially completed blocks left at the end of Part II can be completed in 47 ways. Each of the resulting 4-(12, 6, 4) designs belongs to one of nine equivalence classes.

## 2. Parameters and Block Types for 4 —(12, 6, 4) Designs

For any  $t-(v, k, \lambda)$  design let  $\lambda_i$  be the number of times each  $i$ -subset of the  $v$  points appears in the design. Thus  $\lambda_0 = b$  is the number of blocks;  $\lambda_1 = r$  is the number of replicas of each point; and  $\lambda_t = \lambda$ . Then it is well-known that

$$\lambda_i = \frac{(v-i)(v-i-1)\dots(v-t+1)}{(k-i)(k-i-1)\dots(k-t+1)}\lambda; \quad \text{where } 0 \leq i \leq t.$$

For a 4-(12, 6, 4) design we have

$$\lambda_0 = b = 132, \quad \lambda_1 = r = 66, \quad \lambda_2 = 30, \quad \lambda_3 = 12, \quad \lambda_4 = 4.$$

Let  $S$  be any 6-subset of the 12 points of a 4-(12, 6, 4) design and let  $b_i$  be the number of blocks intersecting  $S$  in exactly  $i$  points. Then by counting successively blocks, point occurrences, pair occurrences, etc, in two ways, we have for the *block intersection numbers* the equations;

$$\begin{aligned} b_0 + b_1 + b_2 + b_3 + b_4 + b_5 + b_6 &= 132, \\ b_1 + 2b_2 + 3b_3 + 4b_4 + 5b_5 + 6b_6 &= 396, \\ 2b_2 + 6b_3 + 12b_4 + 20b_5 + 30b_6 &= 900, \\ 6b_3 + 24b_4 + 60b_5 + 120b_6 &= 1440, \\ 24b_4 + 120b_5 + 360b_6 &= 1440. \end{aligned}$$

From these  $b_0 = 6 - b_5 - 5b_6$ . But  $b_6 \geq 1$  so  $b_6 = 1$  and there are no repeated blocks. Therefore

$$\begin{aligned} b_0 &= 1 - b_5, \quad b_1 = 5b_5, \quad b_2 = 45 - 10b_5, \quad b_3 = 40 + 10b_5, \quad b_4 = 45 - 5b_5, \\ b_5 &= 0 \text{ or } 1; \end{aligned}$$

and only two solution sets are possible. They are :

	$b_0$	$b_1$	$b_2$	$b_3$	$b_4$	$b_5$	$b_6$
Type I	1	0	45	40	45	0	1
Type II	0	5	35	50	40	1	1

The blocks of either type occur in pairs. A block of Type I is disjoint from just one other block. A block of Type II has five points in common with just one other block and intersects all other blocks. Two blocks of Type II with five points in common are said to be *friendly* blocks.

**THEOREM:** *If all the blocks of a 4-(12, 6, 4) design are of Type I then the design is a 5-(12, 6, 1) design.*

**Proof:** No quintuple of points can occur more than once in the design. The 132 blocks of the design each contain  $\binom{6}{5} = 6$  distinct quintuples so the design contains 792 distinct

quintuples in all. But  $\binom{12}{5} = 792$  so each possible quintuple occurs just once and the design must be a 5-(12, 6, 1) design.  $\square$

The design is the well-known 5-(12, 6, 1) design associated with the Mathieu group  $M_{12}$ .

As a by-product of the work above we have a short proof of a result of Dehon[4] and Oberschelp[5].

**THEOREM:** A 4-(12, 6, 2) design cannot exist.

**Proof:** If such a design did exist then, by the duplication of each block, a 4-(12, 6, 4) design with repeated blocks could be made; but no 4-(12, 6, 4) design can have repeated blocks.  $\square$

### 3. The Distribution of the Seven Distinct Points from a Pair of Type II Blocks.

To investigate 4-(12, 6, 4) designs which are not 5-(12, 6, 1) designs it must be assumed that such designs have at least one pair of Type II blocks. In that case there is at least one quintuple of points which occurs twice in the design. We take the pair of blocks to be [123456] and [123457]. The two points, 6 and 7, which do not belong to both of these friendly blocks, play special roles and will be called *prongs*. Thus each pair of Type II blocks has associated with it a *prong pair*, in this case 67. The intention is to show that the seven points associated with a pair of Type II blocks are distributed over the other blocks in a unique pattern. This gives a unique skeleton for a 4-(12, 6, 4) design in which the placing of the seven points 1, 2, 3, 4, 5, 6, 7 is determined. The remaining five points, 8, 9, a, b, c say, then have to be judiciously inserted to complete the design. Towards this end we embark on a series of lemmas.

**LEMMA:** Let  $N_5$  be the number of blocks containing a given quintuple of points. Let  $E_0$  be the number of blocks containing none of the quintuple. Then

$$E_0 = 2 - N_5.$$

**Proof:** By the principle of inclusion and exclusion we have

$$\begin{aligned} E_0 &= b - \binom{5}{1}r + \binom{5}{2}\lambda_2 - \binom{5}{3}\lambda_3 + \binom{5}{4}\lambda_4 - N_5 \\ &= 132 - 5.66 + 10.30 - 10.12 + 5.4 - N_5 \\ &= 2 - N_5. \quad \square \end{aligned}$$

**LEMMA:** Let [123456] and [123457] be a pair of friendly blocks in a 4-(12, 6, 4) design. Then any block intersecting one of the pair in just one point contains the prong of the other.

**Proof:** The quintuple 23456 cannot occur elsewhere in the design. By the previous

lemma,  $E_0 = 2 - N_5 = 1$ , so there is one block not containing any of 2, 3, 4, 5, 6. But the type II block [123456] intersects all other blocks. Therefore there must be a block A, [1 . . . .], not containing any of 2, 3, 4, 5, 6. Now, there are two blocks not containing any of 8, 9, a, b, c, namely the given pair of type II blocks. Therefore, by the previous lemma again, 8, 9, a, b, c cannot all be in the same block; for otherwise  $E_0 \neq 2$ . Hence the block A must contain 7 which is the prong of [123457].  $\square$

LEMMA: *If a 4-(12, 6, 4) design contains the pair of type II blocks [123456] and [123457] then it contains the ten blocks*

[16 . . . .], [26 . . . .], [36 . . . .], [46 . . . .], [56 . . . .],  
[17 . . . .], [27 . . . .], [37 . . . .], [47 . . . .], [57 . . . .],

*in which the dots represent elements from the set {8, 9, a, b, c}.*

Proof: This is a direct consequence of the previous lemma since 1,2,3,4,5 are interchangeable and 6 and 7 are interchangeable.  $\square$

LEMMA: *If a 4-(12, 6, 4) design contains friendly blocks [123456] and [123457] then it contains the ten blocks*

[167 . . .], [267 . . .], [367 . . .], [467 . . .], [567 . . .],  
[167 . . .], [267 . . .], [367 . . .], [467 . . .], [567 . . .],

*in which the dots represent elements from the set {8, 9, a, b, c}.*

Proof: If  $E_0$  is the number of blocks not containing any of the quadruple 2, 3, 4, 5, then by the principle of inclusion and exclusion,

$$E_0 = 132 - \binom{4}{1}r + \binom{4}{2}\lambda_2 - \binom{4}{3}\lambda_3 + \binom{4}{4}\lambda_4 = 132 - 264 + 180 - 48 + 4 = 4.$$

But all the blocks in the design must contain members of the quintuple 1, 2, 3, 4, 5. Therefore the four blocks not containing the quadruple 2, 3, 4, 5, all contain 1. Two of these four blocks are the blocks [16 . . . .], [17 . . . .] of the previous lemma. The remaining two, which must intersect both of [123456] and [123457] in at least two points, must be [167 . . .] and [167 . . .] where the dots represent elements of the set {8, 9, a, b, c}. Cycling through 1, 2, 3, 4, 5 produces the ten blocks in the statement of the lemma.  $\square$

LEMMA: *If a 4-(12, 6, 4) design contains the pair of friendly blocks [123456] and [123457] then the prongs, 6 and 7, cannot be orphans; that is to say, they cannot appear on a block either individually or together unless accompanied by at least one of the non-prong points 1, 2, 3, 4, 5.*

Proof: This is a consequence of the two previous lemmas.  $\square$

#### 4. The Proto-skeleton

Given the pair of friendly blocks [123456] and [123457], it is a matter of simple counting to determine the distribution of the six points 1, 2, 3, 4, 5, 6 over the rest of the blocks in the design. With the help of the previous lemmas some of the 7's can also

be placed. To facilitate further progress, in Table 1 there is presented a *proto-skeleton* in which the two specified friendly blocks are in section A and the other 130 partially

A	C	D	E
123456	1236 .. 1456 ..	126 ... 246 ...	167 ...
123457	1236 .. 1456 ..	126 ... 246 ...	167 ...
	1236 .. 1456 ..	12 ... 24 ...	
	123 ... 145 ...	12 ... 24 ...	267 ...
	123 ... 145 ...		267 ...
	123 ... 145 ...	136 ... 256 ...	
		136 ... 256 ...	367 ...
		13 ... 25 ...	367 ...
		13 ... 25 ...	
	1246 .. 2346 ..	146 ... 346 ...	467 ...
	1246 .. 2346 ..	146 ... 346 ...	467 ...
	1246 .. 2346 ..	14 ... 34 ...	
	124 ... 234 ...	14 ... 34 ...	567 ...
	124 ... 234 ...	14 ... 34 ...	567 ...
	124 ... 234 ...		
	1256 .. 2356 ..	156 ... 356 ...	
	1256 .. 2356 ..	156 ... 356 ...	
	1256 .. 2356 ..	15 ... 35 ...	
	125 ... 235 ...	15 ... 35 ...	
	125 ... 235 ...	15 ... 35 ...	
	125 ... 235 ...		
	1346 .. 2456 ..	236 ... 456 ...	
	1346 .. 2456 ..	236 ... 456 ...	16 ...
	1346 .. 2456 ..	23 ... 45 ...	17 ...
	134 ... 245 ...	23 ... 45 ...	
	134 ... 245 ...		26 ...
	134 ... 245 ...		27 ...
	1356 .. 3456 ..		
	1356 .. 3456 ..		36 ...
	1356 .. 3456 ..		37 ...
	135 ... 345 ...		
	135 ... 345 ...		46 ...
	135 ... 345 ...		47 ...
	135 ... 345 ...		
	135 ... 345 ...		56 ...
			57 ...

Table 1 The Proto-Skeleton

completed blocks are grouped into sections B, C, D, E and F. The blocks of B intersect those of A in four points but do not contain 6 or 7. The blocks of C contain just three of 1, 2, 3, 4, 5 and may contain 6 or 7 or both. The blocks of D contain just two of 1, 2, 3, 4, 5 and may contain 6 or 7 or both. The blocks of E and F contain just one of 1, 2, 3, 4, 5 and are the blocks described in the previous two lemmas. The positions of the 6's within sections C and D are all determined. The placings of the 7's within

these sections are not so easily decided. Within each subsection of section C the point 7 must appear just three times; within each subsection of D the point 7 must appear just twice. These appearances are subject to each quadruple of points appearing just four times in the completed design.

Here we make use of the rule of five (RF for short). This rule, which encapsulates a very basic principle, is used frequently in the subsequent work.

LEMMA: *The rule of five; Given any three blocks of a 4-(12, 6, 4) design, at most two of them can have five points in common.*

Proof: The blocks are of either Type I or of Type II and any one of them can intersect at most one other block of the design in at most five points.  $\square$

An immediate application of the RF proves the following lemma.

LEMMA: *Each subsection of section C of the proto-skeleton can contain the pair 67 at most twice.*  $\square$

LEMMA: *In section C of the proto-skeleton there are exactly 10 blocks containing both 6 and 7.*

Proof: A quadruple from the set  $\{8, 9, a, b, c\}$  can be chosen in five ways. Consider the four points 8, 9, a, b. There are four blocks that contain all of them. The complement of a 4-(12, 6, 4) design is also a 4-(12, 6, 4) design. Therefore there are four blocks that contain none of 8, 9, a, b. Two of these blocks are [123456] and [123457]. The other two blocks must be in section C and must contain three of 1, 2, 3, 4, 5 together with 6, 7, and c.  $\square$

With these constraints a computer search showed that there are twenty non-isomorphic ways of inserting the 7's into sections C and D. The following parity argument ruled out fifteen of these ways.

LEMMA: *Let x be one of 1, 2, 3, 4, 5. Then the triple x67 cannot occur an odd number of times in section C of the proto-skeleton.*

Proof: Each of the five quadruples 167y, where  $y \in \{8, 9, a, b, c\}$ , occurs four times in the design, making twenty such quadruples in all. Six of these occur in the first two blocks of section E. Any block of section D containing 167 contains just two of 8, 9, a, b, c. Any block of section C containing 167 contains just one of 8, 9, a, b, c. Therefore the number of appearances of 167 in section C must be even.  $\square$

One of the five surviving ways contains, in section C, the six-block configuration

[12367 .],	[12467 .],	[13467 .],
[12367 .],	[12467 .],	[13467 .].

Of these blocks, those with 12 require four distinct points from  $\{8, 9, a, b, c\}$  for completion if the RF is to hold. But then the remaining two blocks cannot be completed without breaking the RF or avoiding repeated blocks. Thus the number of non-isomorphic ways of inserting the 7's into sections C and D is reduced to four, as given in the patterns of Table 2. Once the 7's have been placed in section C, the placement of the remaining 7's, in section D, is forced. Therefore in Table 2 the patterns for section C only are

presented. Since section C must contain ten copies of the pair 67 with at most two copies per subsection, and supposing that these copies are in the first two blocks of the subsection, it suffices to give the first two blocks only for each subsection to determine the whole pattern. The four patterns are given in Table 2.

Pattern I	Pattern II	Pattern III	Pattern IV
12367 . 1236 ..	12367 . 12367 .	12367 . 12367 .	12367 . 12367 .
12467 . 1246 ..	12467 . 12467 .	12467 . 1246 ..	12467 . 1246 ..
12567 . 1256 ..	1256 .. 1256 ..	12567 . 1256 ..	1256 .. 1256 ..
13467 . 1346 ..	1346 .. 1346 ..	13467 . 1346 ..	1346 .. 1346 ..
13567 . 1356 ..	13567 . 13567 .	13567 . 1356 ..	13567 . 1356 ..
14567 . 1456 ..	1456 .. 1456 ..	1456 .. 1456 ..	14567 . 14567 .
23467 . 2346 ..	2346 .. 2346 ..	2346 .. 2346 ..	23467 . 2346 ..
23567 . 2356 ..	2356 .. 2356 ..	2356 .. 2356 ..	23567 . 2356 ..
24567 . 2456 ..	24567 . 24567 .	24567 . 24567 .	24567 . 2456 ..
34567 . 3456 ..	34567 . 34567 .	34567 . 34567 .	34567 . 3456 ..

Table 2 The four inequivalent ways of placing the ten 67 pairs in section C. Only the first two blocks of each subsection are given.

## 5. The Skeleton for a 4-(12, 6, 4) Design

The four patterns can be arrived at "by hand," at the peril of overlooking a case. This was the way they were first produced. A reasonably short proof that Pattern II is not viable is available although it is not given here. Before doing the computer run we knew that Pattern I can be completed to a 4-(12, 6, 4) design as we already had examples of such

A	C	D	E
123456	12367 . 14567 .	1267 . . 2467 . .	167 . . .
123457	1236 . . 1456 . .	126 . . . 246 . . .	167 . . .
	1236 . . 1456 . .	127 . . . 247 . . .	
	1237 . . 1457 . .	12 . . . . 24 . . . .	267 . . .
	1237 . . 1457 . .		267 . . .
	123 . . . 145 . . .	1367 . . 2567 . .	
		136 . . . 256 . . .	367 . . .
		137 . . . 257 . . .	367 . . .
		13 . . . . 25 . . . .	
	12467 . 23467 .		467 . . .
	1246 . . 2346 . .	1467 . . 3467 . .	467 . . .
	1246 . . 2346 . .	146 . . . 346 . . .	
	1247 . . 2347 . .	147 . . . 347 . . .	567 . . .
	1247 . . 2347 . .	14 . . . . 34 . . . .	567 . . .
	124 . . . 234 . . .		
		1567 . . 3567 . .	
	12567 . 23567 .	156 . . . 356 . . .	
	1256 . . 2356 . .	157 . . . 357 . . .	
	1256 . . 2356 . .	15 . . . . 35 . . . .	
	1257 . . 2357 . .		
	1257 . . 2357 . .	2367 . . 4567 . .	
	125 . . . 235 . . .	236 . . . 456 . . .	16 . . . .
		237 . . . 457 . . .	17 . . . .
		23 . . . . 45 . . . .	
	13467 . 24567 .		26 . . . .
	1346 . . 2456 . .		27 . . . .
	1346 . . 2456 . .		
	1347 . . 2457 . .		36 . . . .
	1347 . . 2457 . .		37 . . . .
	1347 . . 2457 . .		
	134 . . . 245 . . .		46 . . . .
			47 . . . .
	13567 . 34567 .		56 . . . .
	1356 . . 3456 . .		57 . . . .
	1356 . . 3456 . .		
	1357 . . 3457 . .		
	1357 . . 3457 . .		
	135 . . . 345 . . .		

Table 3 The skeleton for a 4-(12, 6, 4) design. This is also the paradigm for the distribution of the points from a friendly pair of blocks.

designs. None of the known designs seemed to have Patterns II, III and IV embedded in it and endeavours to make designs having these patterns seemed to go nowhere. Therefore an attempt was made to prove that they cannot lead to completed 4-(12, 6, 4) designs. One of us (DRB) has a very long proof (about fifteen pages) that the Patterns II, III and IV are not viable, a proof which is of the into-the-corner-painting type, and which, since it does not involve novel techniques, will not be given here. However, it was fairly

easy to show that Pattern II is not viable; the somewhat asymmetrical Patterns III and IV are the difficult cases. A computer search eliminated all but Pattern I. The program was tested on Pattern II where the result was known beforehand. Then a short run on Patterns III and IV confirmed that no 4–designs can be completed from them.

The combination of Pattern I and the proto-skeleton gives Table 3, the skeleton of a 4–(12, 6, 4) design with a friendly pair of blocks. Note that the seven distinct points from any pair of friendly blocks will generate the same skeleton, which is therefore the paradigm for the distribution of points from any friendly pair of blocks in the design even though they may not be in section B. The blocks in section F must all be of Type II since they intersect other blocks, namely those of section A, in just one point. Also, in section C there are the two blocks [1236 . . ], [1236 . .]. There is only *one* other block in the design that could possibly be disjoint from one of these, and that is in section D, namely, the block [457 . . .]. Therefore section C contains at least twenty Type II blocks and the design contains at least 32 such blocks.

## 6. Acknowledgements

This work has been carried out while two of us, DRB and APS, have been on visits to each other's institutions. We thank both the Mathematics Departments of the Universities of Canterbury and Queensland for the hospitality given to us. The work was sponsored by the Australian Research Council, from which APS holds an Australian Senior Research Fellowship.

## 7. References

- [1]. D. R. Breach, Sylvia H. Elmes, Martin J. Sharry and Anne Penfold Street, *The curious property of the 4–(12, 6, 4) designs*, submitted for publication (1991).
- [2]. D. R. Breach, *In search of 4–(12, 6, 4) designs; Part II*, in preparation.
- [3]. D. R. Breach and Anne Penfold Street, *In search of 4–(12, 6, 4) designs; Part III*, in preparation.
- [4]. M. Dehon, *Non-existence d'un 3–design de paramètres  $\lambda = 2$ ,  $k = 5$  et  $v = 11$* , *Discrete Mathematics* **15**(1976) 23–25.
- [5]. W. Oberschelp, *Lotto-Guarantiesystem und Block-Plane*, *Mathematisch-Phys. Semesterberichte* **19**(1972) 55–67.