

# Antarctic microbial mats: A modern analog for Archean lacustrine oxygen oases

Dawn Y. Sumner<sup>1</sup>, Ian Hawes<sup>2</sup>, Tyler J. Mackey<sup>1</sup>, Anne D. Jungblut<sup>3</sup>, and Peter T. Doran<sup>4</sup>

<sup>1</sup>Department of Earth and Planetary Sciences, University of California–Davis, 1 Shields Avenue, Davis, California 95616, USA

<sup>2</sup>Gateway Antarctica, University of Canterbury, Christchurch 8041, New Zealand

<sup>3</sup>Department of Life Sciences, The Natural History Museum, Cromwell Road, London SW7 5BD, UK

<sup>4</sup>Department of Geology and Geophysics, Louisiana State University, Baton Rouge, Louisiana 70803, USA

## ABSTRACT

The evolution of oxygenic photosynthesis was the most important geochemical event in Earth history, causing the Great Oxidation Event (GOE) ~2.4 b.y. ago. However, evidence is mixed as to whether O<sub>2</sub> production occurred locally as much as 2.8 b.y. ago, creating O<sub>2</sub> oases, or initiated just prior to the GOE. The biogeochemical dynamics of possible O<sub>2</sub> oases have been poorly constrained due to the absence of modern analogs. However, cyanobacteria in microbial mats in a perennially anoxic region of Lake Fryxell, Antarctica, create a 1–2 mm O<sub>2</sub>-containing layer in the upper mat during summer, providing the first known modern analog for formation of benthic O<sub>2</sub> oases. In Lake Fryxell, benthic cyanobacteria are present below the oxycline in the lake. Mat photosynthesis rates were slow due to low photon flux rate (1–2 μmol m<sup>-2</sup> s<sup>-1</sup>) under thick ice cover, but photosynthetic O<sub>2</sub> production was sufficient to sustain up to 50 μmol O<sub>2</sub> L<sup>-1</sup>, sandwiched between anoxic overlying water and anoxic sediments. We hypothesize that Archean cyanobacteria could have similarly created O<sub>2</sub> oases in benthic mats prior to the GOE. Analogous mats may have been at least partly responsible for geological evidence of oxidative weathering prior to the GOE, and habitats such as Lake Fryxell provide natural laboratories where the impact of benthic O<sub>2</sub> oases on biogeochemical signatures can be investigated.

## INTRODUCTION

The most significant geochemical change in Earth's history was caused by oxygenic photosynthesis in cyanobacteria (Farquhar et al., 2011; Kasting, 2013; Kump et al., 2013; Lyons et al., 2014). Prior to ~2.5 b.y. ago, most environments on Earth's surface were anoxic. However, between 2.45 and 2.32 b.y. ago, the Great Oxidation Event (GOE) led to substantial changes in atmospheric and oceanic chemistry, including the accumulation of free molecular oxygen in the atmosphere and shallow oceans, the loss of reduced ions such as Fe(II) and Mn(II) from shallow seawater, and more robust sulfur cycling through multiple oxidation states (Farquhar et al., 2011; Kasting, 2013; Kump et al., 2013; Lyons et al., 2014). Prior to 2.45 Ga, abundant evidence suggests the presence of either local or temporally short oxidative environments, possibly including O<sub>2</sub> (e.g., Kasting, 1992; Eigenbrode and Freeman, 2006; Anbar et al., 2007; Bosak et al., 2009; Duan et al., 2010; Kendall et al., 2010; Czaja et al., 2012). Evidence for local O<sub>2</sub> accumulation before the GOE can be reconciled with an anoxic atmosphere if “oxygen oases” emerged with sufficient oxygenic photosynthetic activity to produce local oxygenated environments, but insufficient O<sub>2</sub> production to cause a global change in oxidation state (e.g., Kasting, 1992; Eigenbrode and Freeman, 2006; Kendall et al., 2010; Olson et al., 2013; Reinhard et al., 2013; Lalonde and Konhauser, 2015).

Oxygen oases have been proposed for open oceans and coastal waters (Fischer, 1965; Kast-

ing, 1992; Olson et al., 2013; Reinhard et al., 2013) and more recently for terrestrial environments (Lalonde and Konhauser, 2015). In the pelagic oceans, rapid mixing and gas exchange with the atmosphere would have made it difficult for more than a few micromoles of O<sub>2</sub> per liter to accumulate in Archean seawater (Olson et al., 2013; Reinhard et al., 2013). In contrast, benthic microbial mats have relatively low exchange rates, and solute fluxes are limited to diffusion through the boundary layer separating mats from the bulk water column. Thus, it is more likely that the first O<sub>2</sub> oases, and those with the highest O<sub>2</sub> concentrations, formed in benthic mats rather than in pelagic environments (e.g., Herman and Kump, 2005; Lalonde and Konhauser, 2015). Terrestrial O<sub>2</sub> oases are of particular interest for understanding geochemical indications of oxidative weathering

prior to 2.5 Ga; sulfide minerals might have oxidized within Archean soil and freshwater cyanobacterial mats without O<sub>2</sub> accumulating in the atmosphere (Reinhard et al., 2013; Lalonde and Konhauser, 2015).

Evaluating the possible extent and weathering potential of Archean freshwater O<sub>2</sub> oases is difficult due to a paucity of modern analogs; none have been previously identified in anoxic environments. Although O<sub>2</sub> concentrations in cyanobacterial mats are commonly higher than in their environment during the day, their biogeochemical dynamics are likely different than in O<sub>2</sub> oases on a reducing Earth due to the prevalence of O<sub>2</sub> rather than reduced ions in the overlying water column. In contrast, benthic mats in Lake Fryxell, McMurdo Dry Valleys, Antarctica, accumulate O<sub>2</sub> during the summer below a poorly mixed, anoxic water column. Here, we describe the conditions under which this localized O<sub>2</sub> oasis forms and consider implications for Archean lacustrine O<sub>2</sub> oases.

## LAKE FRYXELL

Lake Fryxell (75°35'S, 163°35'E; Fig. 1) is a perennially ice-covered lake in the McMurdo Dry Valleys, Antarctica. The lake occupies a closed basin; meltwater streams flow into the lake during summer, but water is lost only through ablation of ice from the surface and water evaporation from a summer “moat” of melt water around the margins of the lake (Lawrence and Hendy, 1985). Historical imbalances in water supply and loss resulted in evaporation and refilling events, creating density stratification of the lake due to increasing salinity with depth (Lyons et al., 2005). This stratification inhibits




Figure 1. Location of Lake Fryxell in Taylor Valley, one of the McMurdo Dry Valleys, Antarctica.

mixing, as does the permanent ice cover; solute transport below 5 m is dominated by diffusion. The perennial ice cover, which was 4–5 m thick in A.D. 2012, affects gas equilibration with the atmosphere. Dissolved gases enter the lake through stream inflow and are excluded from ice as water freezes, leading to gas accumulation in the upper water column over time. Atmospheric gases such as  $N_2$ ,  $O_2$ , and noble gases are present at concentrations well above atmospheric saturation and are prevented from ebullition by hydrostatic pressure (Craig et al., 1992). Vertical gas transport is, however, limited to diffusion, and, while  $O_2$  is present at high concentrations near the surface of the lake, microbial metabolic processes produce an oxycline within the lake at 9–10 m depth (Vincent, 1981). The water column is euxinic immediately below the oxycline (Green et al., 1989).

Irradiance in Lake Fryxell is highly seasonal with about four months of complete winter darkness. During the summer, 0.5% to 3% of incident light penetrates the ice cover, and illumination declines with depth with an extinction coefficient of 0.5–0.6  $m^{-1}$  (Howard-Williams et al., 1998). There is a sharp chlorophyll maximum in the planktonic community at ~8 m, just above the depth at which  $O_2$  concentration and planktonic net photosynthesis fall to zero (e.g., Vincent, 1981; McKnight et al., 2000; Burnett et al., 2006; Vick-Majors et al., 2014). There are extensive benthic mat communities extending from the moat around the margin of the lake to ~11 m depth that are dominated by cyanobacteria, diatoms, and heterotrophic bacteria (e.g., Wharton et al., 1983; Taton et al., 2003).

## METHODS

All measurements were obtained in November 2012. Observations of water column  $O_2$  were made using an  $O_2$  microelectrode (Unisense, www.unisense.com) with an outside tip diameter of 50  $\mu m$  and a 90% response time of <1 s, connected to a Unisense UW-M underwater logging picoammeter. This was mounted alongside a Brancker Instruments Concerto conductivity-temperature depth (CTD) profiler, so that the tip of the  $O_2$  electrode and the sensing element of the CTD were aligned in similar horizontal planes during deployment. These instruments were deployed by a diver along a fixed transit line at regular intervals to obtain dissolved oxygen, temperature, conductivity, and water depth 50 mm above the mat surface.

Incident irradiance to the lake surface was measured at 15 min intervals at a lakeside weather station (Long Term Ecological Research, www.mcmlter.org). Downwelling irradiance at the lake floor was determined by a diver carrying a LiCor Li192 photosynthetically available radiation (PAR) sensor along the transect line and recording irradiance at depths from 8.9 m to 11 m. Simultaneous measurements on

the lake surface allowed the percent of incident irradiance to be calculated for each depth.

The dominant cyanobacterial and diatom morphotypes in different microbial mat sections were identified on site using an Olympus light microscope (BX51) at 400 $\times$ –1000 $\times$  magnification. Cyanobacteria were assigned to genera based on Komárek and Anagnostidis (1989, 2000, 2005; Table DR1 in the [GSA Data Repository](#)<sup>1</sup>) and diatoms identified as completely as possible based on Spaulding et al. (2008).

Profiles of  $O_2$  in mats were measured by divers using the same Clark-type underwater  $O_2$  microelectrode and picoammeter described above. The electrode was mounted in a manually operated micromanipulator on an aluminum post driven into the sediment (Vopel and Hawes, 2006). After a 24 h stabilization period, the diver returned to the stake and used the micromanipulator to move the electrode normal to the mat surface from a position above the boundary layer to a depth of up to 17 mm into the mat in 1.0 mm increments. The position of the mat surface was estimated by the diver and confirmed by a break in the dissolved  $O_2$  versus depth profile. The diffusive flux of  $O_2$  from the microbial mat into the overlying bottom water and the underlying sediments was calculated from the measured steady-state  $O_2$  gradients according to methods described by Vopel and Hawes (2006) and Hawes et al. (2014).


## RESULTS

### Conductivity, Temperature, Oxygen, and Irradiance

Conductivity increased steadily between 6 m and 11 m depth, from <1 to 5  $mS\ cm^{-1}$ , while water temperature ranged from slightly above 2.5  $^{\circ}C$  to slightly less than 2.75  $^{\circ}C$ , demonstrating that increasing salt content caused density stabilization of the water column. The water column was supersaturated with  $O_2$  to a depth of 9.1 m (1 atm saturation at ambient temperature is ~450  $\mu mol\ L^{-1}$ ), transitioning to complete anoxia below ~9.8 m (Fig. 2). We refer to the depth at which  $O_2$  is unmeasurable as the “ $O_2$  limit”.

During November 2012, the average daily PAR flux incident to the Lake Fryxell weather station ranged from 277 to 783 (average = 600)  $\mu mol\ photons\ m^{-2}\ s^{-1}$ . Average daily maxima and minima were 1186 and 96  $\mu mol\ photons\ m^{-2}\ s^{-1}$ , respectively. The percent surface incident PAR

<sup>1</sup>GSA Data Repository item 2015298, Table DR1 (macroscopic characteristics of cyanobacteria morphotypes); Table DR2 (microelectrode profiles of dissolved  $O_2$  in microbial mats from 9.0 m depth); Table DR3 (microelectrode profiles of dissolved  $O_2$  in microbial mats from 9.8 m depth); and Figure DR1 (image of *Phormidium* from 9.8 m depth), is available online at [www.geosociety.org/pubs/ft2015.htm](http://www.geosociety.org/pubs/ft2015.htm), or on request from [editing@geosociety.org](mailto:editing@geosociety.org) or Documents Secretary, GSA, P.O. Box 9140, Boulder, CO 80301, USA.



**Figure 2.** Depth profile of dissolved  $O_2$  in water column of Lake Fryxell, Antarctica. Error bars represent mean and standard deviation of all measurements within a 10 cm depth bin.


reaching the lake floor fell from 0.74% at 8.9 m to 0.27% at the  $O_2$  limit (9.8 m), 0.20% at 10.4 m, and 0.12% at 11.0 m. Combined surface incident and underwater measurements suggest that at the  $O_2$  limit, monthly average PAR was ~1.6  $\mu mol\ photons\ m^{-2}\ s^{-1}$ , which is above the minimum required light flux of ~1  $\mu mol\ photons\ m^{-2}\ s^{-1}$  for oxygenic photosynthesis estimated for nearby Lake Hoare (Hawes et al., 2001, 2014; Vopel and Hawes, 2006), while average maximum daily values exceeded 3  $\mu mol\ photons\ m^{-2}\ s^{-1}$ . The daily average 1  $\mu mol\ photons\ m^{-2}\ s^{-1}$  threshold was reached at 10.4 m depth, well into the anoxic zone of Lake Fryxell.

### Mat Composition

Laminated cohesive microbial mats coated the sediment-water interface to a water depth of more than 10.2 m. Deeper surfaces were still mat covered, but the mat surface was not cohesive. Based on field microscopy, the bulk of the mats at 9.0 m and 9.8 m consisted of cyanobacteria of the *Leptolybya*, *Pseudanabaena*, and *Phormidium* morphotypes (Table DR1). Mats under  $O_2$  supersaturated water at 9.0 m had pinnacles and were dominated by *Leptolybya* morphotypes (<2  $\mu m$  filament width), whereas the 9.8 m mats at the  $O_2$  limit were flat and dominated by a conspicuous green film of a *Phormidium* morphotype with filament diameter of >6  $\mu m$  (Fig. DR1 in the Data Repository; Table DR1), with the diatom *Diadesmis contenta* also present in some samples.

### Dissolved Oxygen Microprofiles and Oxygen Dynamics

Microelectrode  $O_2$  profiles of benthic mats in the oxic part of the lake were similar to those previously seen in the photic zones of similar Antarctic lakes (Fig. 3A; Table DR2; e.g., Hawes et al., 2014). They show an increase in  $O_2$  through the diffusive boundary layer separating the bulk water from the mat, a substantial  $O_2$  subsurface peak of >800  $\mu mol\ O_2\ L^{-1}$  at ~2 mm below the surface of the mat, then a gradual decline in  $O_2$  with depth, and remaining oxic to >17 mm. In contrast, profiles at 9.8 m showed



**Figure 3. Profiles of dissolved oxygen through mat-water interfaces in Lake Fryxell, Antarctica. A: Mean and range of two replicate profiles through a mat in anoxic part of water column (9.0 m depth). B: Mean and one standard deviation of five replicate profiles in anoxic part of water column (9.8 m depth).**

no  $O_2$  in the bulk water column, an increase in  $O_2$  through the few millimeters of water above the mat, and an  $O_2$  peak at  $\sim 1$  mm depth below the mat surface of  $50 \mu\text{mol } O_2 \text{ L}^{-1}$  (10% atmospheric saturation), becoming anoxic again at a mat depth of  $\sim 6$  mm (Fig. 3B; Table DR3).

The spatial distribution of  $O_2$  within the mats allows calculation of net fluxes associated with the  $O_2$  oasis. Diffusion of  $O_2$  away from the  $O_2$  maxima in microbial mats is an indicator of the net rate of photosynthetic  $O_2$  production (Berg et al., 1998; Hawes et al., 2014), and fluxes of  $O_2$  can be calculated using the slopes of  $O_2$  gradients and appropriate diffusion constants. Using a diffusion coefficient ( $D_0$ ) at  $0^\circ\text{C}$  of  $9.13 \times 10^{-6} \text{ cm}^2 \text{ s}^{-1}$  (Broecker and Peng, 1974; modified for temperature according to Li and Gregory, 1974), the flux of  $O_2$  from the 9.8 m mat to the water column was  $\sim 0.04 \mu\text{mol } O_2 \text{ m}^{-2} \text{ s}^{-1}$ . A similar calculation of downward flux suggests that  $0.013 \mu\text{mol } O_2 \text{ m}^{-2} \text{ s}^{-1}$  was diffusing downward into the mat (calculated using a 20% reduction in  $D_0$  within the mat matrix; Vopel and Hawes, 2006). The total export of  $O_2$  from the photic zone was  $0.05 \mu\text{mol } O_2 \text{ m}^{-2} \text{ s}^{-1}$ .

This result is consistent with the expected rate of photosynthesis at the time of  $O_2$  profiling, when irradiance was  $\sim 2.3 \mu\text{mol photons m}^{-2} \text{ s}^{-1}$ . For nearby Lake Hoare, Hawes et al. (2014) presented evidence that microbial mats absorb 50% of incident irradiance into photosynthetic systems, and that the quantum yield of photosynthesis of shade-adapted mats is  $0.06 \text{ mol } O_2 \text{ mol}^{-1} \text{ photons}$ . If the Lake Fryxell mats behave similarly, photosynthetic production by mats at 9.8 m would be  $0.07 \mu\text{mol } O_2 \text{ m}^{-2} \text{ s}^{-1}$ , slightly in

excess of the calculated net flux out of the mat based on the diffusion profile. The excess of  $0.02 \mu\text{mol } O_2 \text{ m}^{-2} \text{ s}^{-1}$  could have been consumed by respiration and sulfide oxidation within the mat.

### DYNAMICS OF OXYGEN OASES

The two key factors required for development of mat  $O_2$  oases are (1)  $O_2$  production that exceeds local consumption, and (2) relatively slow  $O_2$  transport out of the mat. As soon as  $O_2$  production exceeds local consumption,  $O_2$  transiently accumulates. When the export of  $O_2$  to the surrounding environment is sufficiently slow, a stable  $O_2$  oasis develops. One does develop at 9.8 m in Lake Fryxell even at very low photosynthetic rates. In addition, the net annual export of  $O_2$  from mats at 9.8 m and deeper in Lake Fryxell is not sufficient to even seasonally oxidize the local water column, even though photosynthetic  $O_2$  production is sufficient to create the  $O_2$  oasis. The size and temporal persistence of the oasis depends on the spatial distribution and rates of  $O_2$  production and consumption. These rates will change as irradiance fluctuates daily and seasonally. At higher irradiance, a larger  $O_2$  peak and enhanced export to overlying waters and underlying sediments are likely, potentially facilitating oxidation of reduced species such as  $HS^-$ ,  $Fe(II)$ , and  $Mn(II)$  at the boundary between the oasis and anoxic waters. Conversely, at lower irradiance, the mats will contain a smaller  $O_2$  peak, export less  $O_2$ , and support less oxidation of reduced species. During the winter when there is no light, the  $O_2$  oasis is predicted to disappear entirely.

### Implications for Archean Oxygen Oases

We propose that variations in the balance of net  $O_2$  production and flux of reduced species in Lake Fryxell provide a modern analog for development of  $O_2$  oases prior to the GOE. Archean terrestrial aquatic environments as old as 2.7 b.y. commonly contained benthic mats (e.g., Buck, 1980; Buick, 1992; Rye and Holland, 2000). Once these mats contained cyanobacteria, they likely developed  $O_2$  oases even at very low net photosynthetic rates, such as the  $0.05 \mu\text{mol } O_2 \text{ m}^{-2} \text{ s}^{-1}$  observed at 9.8 m in Lake Fryxell. At such low fluxes, all of the  $O_2$  exported from the mat would have been consumed by oxidation of reduced species in the surrounding environment (e.g., Lalonde and Konhauser, 2015). Thus,  $O_2$  oases with tens of micromoles  $O_2$  per liter could have persisted for a long time without oxidizing large habitats. The size, productivity, and frequency of  $O_2$  oases would have gradually increased with the ecological expansion of cyanobacteria and the evolution of more robust oxygenic photosynthesis. Eventually,  $O_2$  oases may have expanded to the open oceans, although  $O_2$  concentrations were likely an order of magnitude lower (Olson et al., 2013; Reinhard et al., 2013). Gradual declines in the concentrations of

reduced species in seawater would have accompanied the spread of oxygenic photosynthesis and decreased the flux of reduced gases to the atmosphere from the oceans. With this expansion, Earth was primed for the GOE.

An accumulation of  $O_2$  in benthic mats can explain some of the geochemical signatures of early weathering. Specifically, the  $\sim 50 \mu\text{mol } O_2 \text{ L}^{-1}$  observed in Lake Fryxell  $O_2$  oases is sufficiently high to allow some pyrite oxidation at reasonable sediment fluxes (Reinhard et al., 2013; Lalonde and Konhauser, 2015). Evidence for oxidative pyrite weathering on land extends back as far as 2.8 b.y. based on models of sulfur fluxes to the oceans (Stüeken et al., 2012) as well as intervals of enhanced molybdenum influx prior to the GOE (Anbar et al., 2007; Duan et al., 2010; Czaja et al., 2012). The “whiffs of oxygen” proposed for these intervals (e.g., Anbar et al., 2007) may record enhanced terrestrial  $O_2$  oasis development rather than changes in the oxidation state of the atmosphere. Using Lake Fryxell  $O_2$  oases as a model, the search for evidence for Archean  $O_2$  can be more precisely targeted at environments where  $O_2$  oases may have had a substantial impact on biogeochemical cycles as well as those environments where the first, small, transient oases may have formed.

### CONCLUSIONS

The presence of transient  $O_2$  oases in Lake Fryxell benthic mats demonstrates that cyanobacteria are capable of producing  $O_2$  oases with sustained concentrations of  $\sim 50 \mu\text{mol } O_2 \text{ L}^{-1} \text{ s}^{-1}$  without oxidizing their environment. These oases provide a model for Archean  $O_2$  oases, which may have formed prior to the oxidation of Earth’s oceans and atmosphere. Similar benthic  $O_2$  oases could have provided environments for oxidative weathering of continental minerals such as pyrite, creating the geochemical signatures indicating “whiffs of oxygen” prior to the GOE.

### ACKNOWLEDGMENTS

The collection of field data was supported by the National Science Foundation Division of Polar Programs through grant number 1115245, the McMurdo Long Term Ecological Research project. Sumner and Mackey were supported by NASA Astrobiology through grant NN13AI60G. We thank both the U.S. Antarctic program and Antarctica NZ for logistic support. We thank James Kasting and an anonymous reviewer for helpful comments.

### REFERENCES CITED

- Anbar, A.D., et al., 2007, A whiff of oxygen before the great oxidation event?: *Science*, v. 317, p. 1903–1906, doi:10.1126/science.1140325.
- Berg, P., Risgaard-Petersen, N., and Rysgaard, S., 1998, Interpretation of measured concentration profiles in sediment pore water: *Limnology and Oceanography*, v. 43, p. 1500–1510, doi: 10.4319/lo.1998.43.7.1500.
- Bosak, T., Liang, B., Sim, S.M., and Petroff, A.P., 2009, Morphological record of oxygenic photosynthesis in conical stromatolites: *Proceedings of the National Academy of Sciences of the*



- United States of America, v. 106, p. 10,939–10,943, doi:10.1073/pnas.0900885106.
- Broecker, W., and Peng, T.-H., 1974, Gas exchange rates between air and sea: *Tellus*, v. 26, p. 185–190, doi:10.3402/tellusa.v26i1-2.9733.
- Buck, S.G., 1980, Stromatolite and ooid deposits within the fluvial and lacustrine sediments of the Precambrian Ventersdorp Supergroup of South Africa: *Precambrian Research*, v. 12, p. 311–330, doi:10.1016/0301-9268(80)90033-9.
- Buick, R., 1992, The antiquity of oxygenic photosynthesis: Evidence from stromatolites in sulphate-deficient Archaean lakes: *Science*, v. 255, p. 74–77, doi:10.1126/science.11536492.
- Burnett, L., Moorhead, D., Hawes, I., and Howard-Williams, C., 2006, Environmental factors associated with deep chlorophyll maxima in Dry Valley lakes, South Victoria Land, Antarctica: *Arctic, Antarctic, and Alpine Research*, v. 38, p. 179–189, doi:10.1657/1523-0430(2006)38[179:EFAWDC]2.0.CO;2.
- Craig, H., Wharton, R.A., and McKay, C.P., 1992, Oxygen supersaturation in ice-covered Antarctic lakes: Biological versus physical contributions: *Science*, v. 255, p. 318–321, doi:10.1126/science.11539819.
- Czaja, A.D., Johnson, C.M., Roden, E.E., Beard, B.L., Voegelin, A.R., and Nagler, T.F., 2012, Evidence for free oxygen in the Neoproterozoic ocean based on coupled iron–molybdenum isotope fractionation: *Geochimica et Cosmochimica Acta*, v. 86, p. 118–137, doi:10.1016/j.gca.2012.03.007.
- Duan, Y., Anbar, A.D., Arnold, G.L., Lyons, T.W., Gordon, G.W., and Kendall, B., 2010, Molybdenum isotope evidence for mild environmental oxygenation before the Great Oxidation Event: *Geochimica et Cosmochimica Acta*, v. 74, p. 6655–6668, doi:10.1016/j.gca.2010.08.035.
- Eigenbrode, J.L., and Freeman, K.H., 2006, Late Archaean rise of aerobic microbial ecosystems: Proceedings of the National Academy of Sciences of the United States of America, v. 103, p. 15,759–15,764, doi:10.1073/pnas.0607540103.
- Farquhar, J., Zerkle, A.L., and Bekker, A., 2011, Geological constraints on the origin of oxygenic photosynthesis: *Photosynthesis Research*, v. 107, p. 11–36, doi:10.1007/s11220-010-9594-0.
- Fischer, A., 1965, Fossils, early life, and atmospheric history: Proceedings of the National Academy of Sciences of the United States of America, v. 53, p. 1205–1215, doi:10.1073/pnas.53.6.1205.
- Green, W.J., Gardner, T.J., Ferdelman, T.F., Angle, M.P., Varner, L.C., and Nixon, P., 1989, Geochemical processes in the Lake Fryxell Basin (Victoria Land, Antarctica): *Hydrobiologia*, v. 172, p. 129–148, doi:10.1007/BF00031617.
- Hawes, I., Moorhead, D., Sutherland, D., Schmeling, J., and Schwarz, A.-M., 2001, Benthic primary production in two perennially ice-covered Antarctic lakes: Patterns of biomass accumulation with a model of community metabolism: *Antarctic Science*, v. 13, p. 18–27, doi:10.1017/S0954102001000049.
- Hawes, I., Giles, H., and Doran, P.T., 2014, Estimating photosynthetic activity in microbial mats in an ice-covered Antarctic lake using automated oxygen microelectrode profiling and variable chlorophyll fluorescence: *Limnology and Oceanography*, v. 59, p. 674–688.
- Herman, E.K., and Kump, L.R., 2005, Biogeochemistry of microbial mats under Precambrian environmental conditions: A modeling study: *Geobiology*, v. 3, p. 77–92, doi:10.1111/j.1472-4669.2005.00048.x.
- Howard-Williams, C., Schwarz, A.-M., Hawes, I., and Priscu, J.C., 1998, Optical properties of the McMurdo Dry Valley lakes, in Priscu, J.C., ed., *Ecosystem Dynamics in a Polar Desert: The McMurdo Dry Valleys, Antarctica*: Washington, D.C., American Geophysical Union, p. 189–203, doi:10.1029/AR072p0189.
- Kasting, J.F., 1992, Models relating to Proterozoic atmospheric and ocean chemistry, in Schopf, J., and Klein, C., eds., *The Proterozoic Biosphere: A Multidisciplinary Study*: Cambridge, UK, Cambridge University Press, p. 1185–1187.
- Kasting, J.F., 2013, What caused the rise of atmospheric O<sub>2</sub>? : *Chemical Geology*, v. 362, p. 13–25, doi:10.1016/j.chemgeo.2013.05.039.
- Kendall, B., Reinhard, C.T., Lyons, T.W., Kaufman, A.J., Poulton, S.W., and Anbar, A.D., 2010, Pervasive oxygenation along late Archaean ocean margins: *Nature Geoscience*, v. 3, p. 647–652, doi:10.1038/ngeo942.
- Komárek, J., and Anagnostidis, K., 1989, Modern approach to the classification system of cyanophytes: 4. Nostocales: *Archiv fuer Hydrobiologie*, v. 82, Supplement, p. 247–345.
- Komárek, J., and Anagnostidis, K., 2000, Cyanoprokaryota: Teil/Part 1. Chroococcales, in Ettl, H., et al., eds., *Süßwasserflora von Mitteleuropa*: Jena, Germany, Gustav Fischer Verlag, p. 1–548.
- Komárek, J., and Anagnostidis, K., 2005, Cyanoprokaryota: Teil/Part 2. Oscillatoriales, in Budek, B., et al., eds., *Süßwasserflora von Mitteleuropa*: Jena, Germany, Gustav Fischer Verlag, p. 1–758.
- Kump, L.R., Fallick, A.E., Melezhik, V.A., Strauss, H., and Lepland, A., 2013, The Great Oxidation Event, in Melezhik, V.A., et al., eds., *Reading the Archive of Earth's Oxygenation, Volume 3: Global Events and the Fennoscandian Arctic Russia–Drilling Early Earth Project*: Berlin, Springer-Verlag, p. 1517–1533.
- Lalonde, S.V., and Konhauser, K.O., 2015, Benthic perspective on Earth's oldest evidence for oxygenic photosynthesis: Proceedings of the National Academy of Sciences of the United States of America, v. 112, p. 995–1000, doi:10.1073/pnas.1415718112.
- Lawrence, M.J.F., and Hendy, C.H., 1985, Water column and sediment characteristics of Lake Fryxell, Taylor Valley, Antarctica: *New Zealand Journal of Geology and Geophysics*, v. 28, p. 543–552, doi:10.1080/00288306.1985.10421206.
- Li, Y.-H., and Gregory, S., 1974, Diffusion of ions in sea water and in deep-sea sediments: *Geochimica et Cosmochimica Acta*, v. 38, p. 703–714, doi:10.1016/0016-7037(74)90145-8.
- Lyons, W.B., Welch, K.A., Snyder, G., Olesik, J., Graham, E.Y., Marion, G.M., and Poreda, R.J., 2005, Halogen geochemistry of the McMurdo Dry Valleys lakes, Antarctica: Clues to the origin of solutes and lake evolution: *Geochimica et Cosmochimica Acta*, v. 69, p. 305–323, doi:10.1016/j.gca.2004.06.040.
- Lyons, T.W., Reinhard, C.T., and Planavsky, N.J., 2014, The rise of oxygen in Earth's early ocean and atmosphere: *Nature*, v. 506, p. 307–315, doi:10.1038/nature13068.
- McKnight, D.M., Howes, B.L., Taylor, C.D., and Goehring, D.D., 2000, Phytoplankton dynamics in a stably stratified Antarctic lake during winter darkness: *Journal of Phycology*, v. 36, p. 852–861, doi:10.1046/j.1529-8817.2000.00031.x.
- Olson, S.L., Kump, L.R., and Kasting, J.F., 2013, Quantifying the areal extent and dissolved oxygen concentrations of Archaean oxygen oases: *Chemical Geology*, v. 362, p. 35–43, doi:10.1016/j.chemgeo.2013.08.012.
- Reinhard, C.T., Lalonde, S.V., and Lyons, T.W., 2013, Oxidative sulfide dissolution on the early Earth: *Chemical Geology*, v. 362, p. 44–55, doi:10.1016/j.chemgeo.2013.10.006.
- Rye, R., and Holland, H.D., 2000, Life associated with a 2.76 Ga ephemeral pond?: Evidence from Mount Roe #2 paleosol: *Geology*, v. 28, p. 483–486, doi:10.1130/0091-7613(2000)28<483:LAWAGE>2.0.CO;2.
- Spaulding, S., et al., 2008, Antarctic freshwater diatoms: McMurdo Dry Valleys Long Term Ecological Research: Boulder, Colorado, <http://huey.colorado.edu/diatoms> (accessed August 2015).
- Stüeken, E.E., Catling, D.C., and Buick, R., 2012, Contributions to late Archaean sulphur cycling by life on land: *Nature Geoscience*, v. 5, p. 722–725, doi:10.1038/ngeo1585.
- Taton, A., Grubisic, S., Brambilla, E., De Wit, R., and Wilimotte, A., 2003, Cyanobacterial diversity in natural and artificial microbial mats of Lake Fryxell (McMurdo Dry Valleys, Antarctica): A morphological and molecular approach: *Applied and Environmental Microbiology*, v. 69, p. 5157–5169, doi:10.1128/AEM.69.9.5157-5169.2003.
- Vick-Majors, T.J., Priscu, J.C., and Amaral-Zettler, L.A., 2014, Modular community structure suggests metabolic plasticity during the transition to polar night in ice-covered Antarctic lakes: *The ISME Journal*, v. 8, p. 778–789, doi:10.1038/ismej.2013.190.
- Vincent, W.F., 1981, Production strategies in Antarctic inland waters: Phytoplankton eco-physiology in a permanently ice-covered lake: *Ecology*, v. 62, p. 1215–1224, doi:10.2307/1937286.
- Vopel, K., and Hawes, I., 2006, Photosynthetic performance of benthic microbial mats in Lake Hoare, Antarctica: *Limnology and Oceanography*, v. 51, p. 1801–1812, doi:10.4319/lso.2006.51.4.1801.
- Wharton, B.C., Jr., Parker, B.C., and Simmons, G.M., Jr., 1983, Distribution, species composition and morphology of algal mats in Antarctic dry valley lakes: *Phycologia*, v. 22, p. 355–365, doi:10.2216/i0031-8884-22-4-355.1.

Manuscript received 13 May 2015

Revised manuscript received 25 July 2015

Manuscript accepted 27 July 2015

Printed in USA