

## A novel maize-infecting mastrevirus from La Réunion Island

Daniel Pande · Simona Kraberger · Pierre Lefeuvre ·  
Jean-Michel Lett · Dionne N. Shepherd ·  
Arvind Varsani · Darren P. Martin

Received: 13 February 2012 / Accepted: 18 March 2012 / Published online: 28 April 2012  
© Springer-Verlag 2012

**Abstract** Despite extensive sampling, only one virus belonging to the genus *Mastrevirus* of the family *Geminiviridae*, maize streak virus (MSV), has until now been detected in maize with maize streak disease (MSD) symptoms. Here, we report for the first time a second, highly divergent, mastrevirus isolated from two maize plants displaying characteristic MSD-like symptoms, sampled on the South-west Indian Ocean Island, La Réunion. The two isolates shared <57 % genome-wide identity with all other known mastreviruses. We propose calling the new species Maize streak Réunion virus.

The genus *Mastrevirus* of the family *Geminiviridae* contains species with circular single-stranded DNA (ssDNA) genomes of approximately 2.7 kb. Mastreviruses infect a wide variety of monocotyledonous and dicotyledonous hosts and are transmitted by leafhopper vectors. Monocot-infecting mastreviruses have previously been found throughout Africa (including the South-west Indian Ocean islands: SWIO), Europe, Japan, Southeast Asia, the island of Vanuatu, and Australia. Mastreviruses previously found on SWIO islands such as La Réunion and Mauritius include various maize streak virus (MSV) strains, sugarcane streak Réunion virus (SSRV) and sugarcane streak virus (SSV) [13, 19, 25, 26].

Here, we report the genome sequences of two isolates (PR50 and PR52) belonging to a divergent mastrevirus species from La Réunion. Leaves were sampled in Saint Pierre (21.6161 S, 55.45784 E and 21.3206 S, 55.48536 E) from two maize plants displaying chlorotic streaks along leaf veins similar to those seen in MSV infections of maize.

GenBank accession numbers: MSRV-[RE-StP-PR52-2009]: JQ624879. MSRV-[RE-StP-PR50-2009]: JQ624880

**Electronic supplementary material** The online version of this article (doi:10.1007/s00705-012-1314-y) contains supplementary material, which is available to authorized users.

D. Pande  
Department of Botany and Horticulture, Maseno University,  
P.O. Box 333, Maseno, Kenya

S. Kraberger · A. Varsani (✉)  
School of Biological Sciences, University of Canterbury, Private  
Bag 4800, Christchurch 8140, New Zealand  
e-mail: arvind.varsani@canterbury.ac.nz

P. Lefeuvre · J.-M. Lett  
CIRAD, UMR 53 PVBMT CIRAD-Université de la Réunion,  
Pôle de Protection des Plantes, Ligne Paradis, 97410 Saint  
Pierre, La Réunion, France

D. N. Shepherd  
Department of Molecular and Cell Biology, University of Cape  
Town, Rondebosch, Cape Town 7701, South Africa

A. Varsani  
Biomolecular Interaction Centre, University of Canterbury,  
Private Bag 4800, Christchurch 8140, New Zealand

A. Varsani  
Electron Microscope Unit, University of Cape Town,  
Rondebosch, Cape Town 7701, South Africa


D. P. Martin  
Computational Biology Group, Institute of Infectious Diseases  
and Molecular Medicine, University of Cape Town, Observatory,  
Cape Town 7925, South Africa

Circular viral DNA molecules were amplified from a crude total DNA extract from the two maize samples using Phi29 DNA polymerase (TempliPhi™, GE Healthcare, USA) as described previously [15, 16, 18] and cut with *KpnI* (PR50) and *BamHI* (PR52) to yield ~2.7-kb DNA fragments, which were subsequently ligated to the *KpnI* and *BamHI* sites of pGEMZf+ (Promega Biotech, USA). The clones were fully sequenced by primer walking at MacroGen Inc. (Korea).


The two viral isolates share 97.3 % nucleotide sequence identity. A NCBI BLASTn [28] analysis of the assembled 2882-bp sequence of the virus isolates from both plants indicated that they shared significant degrees of similarity with mastreviruses in general and in particular with panicum streak virus (PanSV) isolates in the capsid protein (CP) and replication-associated protein (Rep) open reading frames (ORFs; maximum identity of 72 % with 30 % genome-wide coverage E value  $6 \times 10^{-49}$  to PanSV-A [ZM-Nya-g180-2007], GenBank # EU224263). The novel viral genomes were aligned using MUSCLE [6] with representative sequences of mastrevirus species and major strain groupings using MEGA version 5 [24]. Pairwise

**Fig. 2** Maximum-likelihood phylogenetic tree of Rep (A), and CP (B) proteins (constructed using the LG amino acid substitution model) along with the pairwise distance matrix of Rep and CP. Branches marked with filled and open circles were supported in >90 % and 60–89 % of bootstrap replicates respectively, and branches with <50 % bootstrap support have been collapsed. Tree branches are coloured according to the geographical origins of the viruses. *Eragrostis curvula* streak virus (ECSV) is included in both trees despite not being a mastrevirus because it has a very mastrevirus-like capsid protein gene. Beet curly top Iran virus was used to root both the trees


identity analysis (Hamming- or p-distance with pairwise deletion of gaps) indicated that, with the exception of *Eragrostis minor* streak virus (EMSV), the novel viral isolates shared only 53–57 % genome-wide identity with all other known African streak viruses. The novel maize viruses shared only 48.7 % genome-wide identity with EMSV (Fig. 1). It is clear that these new viral genomes represent a new mastrevirus species (based on the currently accepted International Committee on Taxonomy of Viruses 75 % identity mastrevirus species demarcation threshold [4, 7]), and we propose the name Maize streak Réunion virus (MSRV) (Fig. 2).



Branches marked with filled and open circles were supported in >90 % and 60–89 % of bootstrap replicates respectively, and branches with <50 % bootstrap support have been collapsed. The phylogenetic tree is rooted using the full genome sequence of *Eragrostis curvula* streak virus (ECSV; not shown)



**C**



Recombination analysis using RDP3 (with default settings; [12]) failed to reveal any evidence that the virus was recombinant. A maximum-likelihood (ML) phylogenetic tree was therefore constructed from the full alignment using PHYML (version 3 with GTR+I+G<sub>4</sub> chosen as the best-fit model by RDP; [9, 12]; Fig. 1). The ML tree clearly indicated that the new virus is a mastrevirus that is most closely related to the virus clade including *Digitaria* streak virus (DSV) from Vanuatu and the various known African streak viruses (with the notable exception of EMSV, which is more closely related to the Asian streak virus *Miscanthus* streak virus from Japan [5]).

As with almost all known geminiviruses, MSRV contains the nonanucleotide sequence TAATATTAC, which marks its origin of virion-strand DNA replication. The MSRV genomes contain (1) two probable intergenic regions (one short and one long), (2) two probable virion-strand genes (likely encoding a movement protein [MP] and a CP), and (3) an intron-containing complementary-sense gene (likely expressing both a replication-associated protein (Rep) from the spliced transcript and a Rep derivative, RepA, from the unspliced transcript (Supplementary Fig. 1). Various nucleotide sequence motifs that are probably functional within the full-genome nucleotide and ORF sequences of MSRV are shown in Supplementary Figs. 1 and 2, respectively.

While the MP of MSRV shares the highest amino acid sequence identity (42.9 %) with chickpea chlorosis virus B (CpCV-B-[AU:Qld21:2003]-GU256531), its CP is most similar to that of sugarcane streak Egypt virus (60.6 %; SSEV; SSEV-[EG:Egypt:2000]-AF239159). Since it is known that the African streak viruses MSV, PanSV, SSRV, SSEV and SSV are all transmitted by *Cicadulina* species [1–3, 17, 20–23], it is likely that the other viruses in the main African streak virus group are also transmitted by the same species. It is, however, not certain that MSRV is also transmissible by *Cicadulina* species since the CP is the determinant of geminivirus transmission specificity, and the MRSV CP sequence branches basal to the African streak virus group sequences.

The Rep of MSRV shares 56.3 % amino acid identity with MSV-I (MSV-I [ZA-NewA-g217-2007]-EU62863) and 47–55 % identity with those of other African streak virus Reps. Within the MSRV Rep we identified various amino acid motifs previously identified in other mastreviruses including (1) a likely rolling-circle replication (RCR) motif I sequence (FLTYP [11]), (2) a possible metal ion coordination domain (HLHVLLQ, corresponding to RCR motif II [11]), (3) a possible GRS motif (YHPNIQASR [14]), (4) a possible RCR motif III sequence (YILKSP [11]), (5) a probable dNTP-binding domain sequence (VGX<sub>4</sub>GKTTW<sub>30</sub>DD [8]), (6) a retinoblastoma-related protein binding motif (LHCYE [27]), and (7) a potential

oligomerisation domain (SAERLFPTLPSPFV [10]; Supplementary Fig. 2).

MSRV is particularly interesting because despite extensive sampling and analysis of maize displaying maize streak disease symptoms, to date, the only other mastrevirus ever sampled from maize is MSV. Given that maize is the most important food crop in Africa and the possibility that this new virus might already be circulating throughout Africa or that it could be introduced to the continent at some time in the future, it is imperative that we actively pursue the biological characterization of MSRV. Just as members of multiple geminivirus species form a disease complex affecting Africa's other important food crop, cassava, it remains a possibility that MRSV and MSV might together form a disease complex that has a far-reaching impact on African maize production. We are currently in the process of obtaining infectious MSRV clones, which will be used to determine both the capacity of this virus to infect different maize genotypes and whether, as is the case with other African streak viruses, it is transmissible by *Cicadulina* sp.

**Acknowledgments** This work was funded by a grant from the National Foundation of Science of South Africa and a University of Cape Town Block Grant awarded to AV. PL is funded by GIS CRVOI through the EmerGe grant (N°PRAO/AIRD/CRVOI/08/03).

## References

1. Bigarre L, Salah M, Granier M, Frutos R, Thouvenel JC, Peterschmitt M (1999) Nucleotide sequence evidence for three distinct sugarcane streak mastreviruses. *Arch Virol* 144:2331–2344
2. Briddon RW, Lunness P, Chamberlin LC, Pinner MS, Brundish H, Markham PG (1992) The nucleotide sequence of an infectious insect-transmissible clone of the geminivirus *Panicum* streak virus. *J Gen Virol* 73(Pt 5):1041–1047
3. Briddon RW, Lunness P, Bedford ID, Chamberlin LCL, Mesfin T, Markham PG (1996) A streak disease of pearl millet caused by a leafhopper-transmitted geminivirus. *Eur J Plant Pathol* 102:397–400
4. Brown JK, Fauquet CM, Briddon RW, Zerbini M, Moriones E, Navas-Castillo J (2011) Geminiviridae. In: King AMQ, Adams MJ, Carstens EB, Lefkowitz EJ (eds) *Virus taxonomy: classification and nomenclature of viruses: Ninth Report of the International Committee on Taxonomy of Viruses*. Elsevier Academic Press, San Diego, pp 251–374
5. Chatani M, Matsumoto Y, Mizuta H, Ikegami M, Boulton MI, Davies JW (1991) The nucleotide-sequence and genome structure of the Geminivirus *Miscanthus* streak virus. *J Gen Virol* 72:2325–2331
6. Edgar RC (2004) MUSCLE: multiple sequence alignment with high accuracy and high throughput. *Nucleic Acids Res* 32:1792–1797
7. Fauquet CM, Briddon RW, Brown JK, Moriones E, Stanley J, Zerbini M, Zhou X (2008) Geminivirus strain demarcation and nomenclature. *Arch Virol* 153:783–821
8. Gorbalenya AE, Koonin EV (1989) Viral proteins containing the purine NTP-binding sequence pattern. *Nucleic Acids Res* 17:8413–8440

9. Guindon S, Gascuel O (2003) A simple, fast, and accurate algorithm to estimate large phylogenies by maximum likelihood. *Syst Biol* 52:696–704
10. Horvath GV, Pettko-Szandtner A, Nikovics K, Bilgin M, Boulton M, Davies JW, Gutierrez C, Dudits D (1998) Prediction of functional regions of the maize streak virus replication-associated proteins by protein-protein interaction analysis. *Plant Mol Biol* 38:699–712
11. Koonin EV, Ilyina TV (1992) Geminivirus replication proteins are related to prokaryotic plasmid rolling circle DNA-replication Initiator proteins. *J Gen Virol* 73:2763–2766
12. Martin DP, Lemey P, Lott M, Moulton V, Posada D, Lefevre P (2010) RDP3: a flexible and fast computer program for analyzing recombination. *Bioinformatics* 26:2462–2463
13. Monjane AL, Harkins GW, Martin DP, Lemey P, Lefevre P, Shepherd DN, Oluwafemi S, Simuyandi M, Zinga I, Komba EK, Lakoutene DP, Mandakombo N, Mboukoulida J, Semballa S, Tagne A, Tiendrebeogo F, Erdmann JB, van Antwerpen T, Owor BE, Flett B, Ramusi M, Windram OP, Syed R, Lett JM, Briddon RW, Markham PG, Rybicki EP, Varsani A (2011) Reconstructing the history of maize streak virus strain a dispersal to reveal diversification hot spots and its origin in southern Africa. *J Virol* 85:9623–9636
14. Nash TE, Dallas MB, Reyes MI, Buhrman GK, Ascencio-Ibanez JT, Hanley-Bowdoin L (2011) Functional analysis of a novel motif conserved across geminivirus Rep proteins. *J Virol* 85:1182–1192
15. Owor BE, Martin DP, Shepherd DN, Edema R, Monjane AL, Rybicki EP, Thomson JA, Varsani A (2007) Genetic analysis of maize streak virus isolates from Uganda reveals widespread distribution of a recombinant variant. *J Gen Virol* 88:3154–3165
16. Owor BE, Shepherd DN, Taylor NJ, Edema R, Monjane AL, Thomson JA, Martin DP, Varsani A (2007) Successful application of FTA((R)) Classic Card technology and use of bacteriophage phi 29 DNA polymerase for large-scale field sampling and cloning of complete maize streak virus genomes. *J Virol Methods* 140:100–105
17. Peterschmitt M, Reynaud B, Sommermeyer G, Baudin P (1991) Characterization of Maize streak virus isolates using monoclonal and polyclonal antibodies and by transmission to a few hosts. *Plant Dis* 75:27–32
18. Shepherd DN, Martin DP, Lefevre P, Monjane AL, Owor BE, Rybicki EP, Varsani A (2008) A protocol for the rapid isolation of full geminivirus genomes from dried plant tissue. *J Virol Methods* 149:97–102
19. Shepherd DN, Varsani A, Windram OP, Lefevre P, Monjane AL, Owor BE, Martin DP (2008) Novel sugarcane streak and sugarcane streak Reunion mastreviruses from southern Africa and La Reunion. *Arch Virol* 153:605–609
20. Shepherd DN, Martin DP, van der Walt E, Dent K, Varsani A, Rybicki EP (2010) Maize streak virus: an old and complex ‘emerging’ pathogen. *Mol Plant Pathol* 11:1–12
21. Storey HH (1924) The transmission of a new plant virus disease by insects. *Nature* 114:245
22. Storey HH (1925) The transmission of streak disease of maize by the leafhopper *Balclutha mbila* Naude. *Ann Appl Biol* 12:422–439
23. Storey HH, McClean APD (1930) The transmission of streak disease between maize, sugar cane and wild grasses. *Ann Appl Biol* 17:691–719
24. Tamura K, Peterson D, Peterson N, Stecher G, Nei M, Kumar S (2011) MEGA5: Molecular evolutionary genetics analysis using maximum likelihood, evolutionary distance, and maximum parsimony methods. *Mol Biol Evol* 28:2731–2739
25. Varsani A, Oluwafemi S, Windram OP, Shepherd DN, Monjane AL, Owor BE, Rybicki EP, Lefevre P, Martin DP (2008) Panicum streak virus diversity is similar to that observed for maize streak virus. *Arch Virol* 153:601–604
26. Varsani A, Shepherd DN, Monjane AL, Owor BE, Erdmann JB, Rybicki EP, Peterschmitt M, Briddon RW, Markham PG, Oluwafemi S, Windram OP, Lefevre P, Lett JM, Martin DP (2008) Recombination, decreased host specificity and increased mobility may have driven the emergence of maize streak virus as an agricultural pathogen. *J Gen Virol* 89:2063–2074
27. Xie Q, Suarez-Lopez P, Gutierrez C (1995) Identification and analysis of a retinoblastoma binding motif in the replication protein of a plant DNA virus: requirement for efficient viral DNA replication. *EMBO J* 14:4073–4082
28. Zhang Z, Schwartz S, Wagner L, Miller W (2000) A greedy algorithm for aligning DNA sequences. *J Comput Biol* 7:203–214